

6 OCT 5-1959
Cont. Copy

SCIENCE


2 October 1959

AMERICAN ASSOCIATION FOR THE ADVANCEMENT OF SCIENCE

518515



*the
laboratory
microscope
which
fulfills
research
requirements*



For photomicrography and simultaneous observation, binocular photo tube "FS" is available at an additional charge of \$50.00.

the new **LEITZ LABOLUX IIIa**

The Leitz Labolux IIIa is a new laboratory microscope with built-in illumination and the famous Labolux ball-bearing focusing control, which combines both coarse and fine focusing in a single knob. Available with a wide variety of accessories, the Labolux IIIa is recommended for all routine laboratory work and, in addition, can be equipped to fulfill the most exacting research.

A variety of interchangeable tubes is available: monocular, binocular or trinocular (binocular viewing plus photo tube for photomicrography). Tubes can be rotated 360° so that the Labolux IIIa may be faced away from the observer, for increased accessibility to all controls and to the object stage, and to make "conference-viewing" by two consultants more convenient.

Among the condensers available are the Abbe type, the Berek 2-diaphragm condenser, and condensers for phase contrast and dark field observations. The Labolux IIIa is readily adapted to fluorescence microscopy by addition of the Leitz fluorescence accessories. The large stand, in a new contemporary design, is constructed for a lifetime of use with fatigue-free operation and precision performance. All controls, including the knobs for the mechanical stage, are in a low convenient position. High-power objectives have spring-loaded mounts for prevention of damage to lenses and slides.

LABOLUX IIIa, Model S 25/95

- inclined binocular tube S with knurled knob to adjust for proper interpupillary distance • built-in mechanical stage #25 • two-lens condenser #95 • substage unit with rack and pinion focusing accepts sleeve-type condensers • quadruple nosepiece with achromats 3.5X, 10X, 45X and 100X oil immersion, the last two having spring-loaded mounts • paired 10X wide-field eyepieces • horizontal carrying case • 3-step transformer, 6 V, 2.5 Amp. **\$734.00**

For literature and/or a personal demonstration in your laboratory, write: Dept. SC-10



E. LEITZ, INC., 468 FOURTH AVENUE, NEW YORK 16, N.Y.
Distributors of the world-famous products of
Ernst Leitz G. m. b. H., Wetzlar, Germany—Ernst Leitz Canada Ltd.
LEICA CAMERAS • LENSES • MICROSCOPES • BINOCULARS

32959

Q1
S35
2d set

IMPERFECT COPY

What!

4

77

58

every order shipped within 8 hours?

Sounds incredible, but it's true! We at NBCo know the importance of TIME to research organizations. Consequently we have developed our order processing, filling and shipping techniques to a precise science, which requires every order to be shipped within 8 hours of receipt . . . and we can even process emergency shipments within ONE hour! In addition to offering the FASTEST service of any research biochemical house, we are equally proud of our reputation for QUALITY . . . the highest possible quality . . . at the lowest possible price. We invite you to join the thousands of organizations who rely on NBCo . . . the world's largest, most complete research biochemical company.



Our stock of more than 2,500 items includes:

- Over 300 Amino Acids
- Over 90 Peptides
- More than 200 Nucleoproteins, Purines, Pyrimidines
- Miscellaneous Biochemicals
- Vitamins
- Enzymes-Crystalline, Purified
- Growth Factors
- Steroid Hormones
- Fatty Acids
- Antibiotics
- Carbohydrates
- Purified Proteins
- Alkaloids
- Biological Salt Mixture
- Biological Test Material
- Glandular Substances



OUR NEW JULY 1959 CATALOG

containing more than 2,500 items is now ready. Fill out coupon below and mail today for your free copy.

Name _____

Firm or Organization _____

Address _____

City _____ Zone _____ State _____

**NUTRITIONAL
BIOCHEMICALS
CORPORATION**

21012 MILES AVENUE
CLEVELAND 28, OHIO

SCIENCE is published weekly by the AAAS, 1515 Massachusetts Ave., NW, Washington 5, D.C. Entered at the Washington, D.C., Post Office as second class matter under the act of 3 March 1879. Annual subscriptions: \$8.50; foreign postage, \$1.50; Canadian postage, 75¢.

NEW...

SARGENT ANALYTICAL OVEN



Designed & Manufactured by
E. H. Sargent & Co.



- HEAT TRANSFER BY RADIATION AND CONDUCTION
- OVER FOUR SQUARE FEET OF WORKING AREA
- DIRECT READING, SINGLE TURN TEMPERATURE ADJUSTING DIAL
- OPERATING RANGE, 35° TO 200°C
- ALL INSIDE METAL AREAS AND AIR VOLUME COINCIDENT IN TEMPERATURE WITHIN $\pm 2\frac{1}{2}^{\circ}\text{C}$ WHETHER UNLOADED OR FULLY LOADED
- TEMPERATURE AT ANY POINT CONSTANT WITHIN $\pm \frac{1}{2}^{\circ}\text{C}$ WHETHER UNLOADED OR FULLY LOADED

S-64080 OVEN—Analytical, Radiation-Conduction Type, Double Wall, 200°C, Sargent. A new laboratory analytical oven designed to emphasize conduction and radiation as modes of heat transfer and to minimize the contribution of convection, thereby achieving an unprecedented uniformity of temperature throughout the entire cubic space of the isolated aluminum interior chamber.

The inner chamber is obstruction free. Three removable flat aluminum shelves and a flat bottom area provide over four square feet of useable space for moisture samples.

Heat is applied at a very low gradient and over a wide area from a nickel-chromium ribbon element in woven glass sleeving. $2\frac{1}{2}$ inches of fiber glass insulation results in low wattage requirement and permits temperature to be regulated by a special Sargent heavy duty, bimetal regulator.

Temperature is adjusted by simple rotation of a single turn disk on the lower panel. A dial type thermometer is mounted on the door.

Constructed of heavy gauge, welded steel, finished in baked enamel. Inlet and outlet vents are provided. The oven door automatically opens if pressure builds up inside.

Range, 35° to 200°C; inside width, 14 inches; inside height, 13 inches; inside depth, 12 inches; outside width, 19 inches; outside height, $22\frac{1}{2}$ inches; outside depth, 16 inches; net weight, 60 lbs. Complete with three shelves and thermometer. For operation from 115 volt, 50 or 60 cycle A.C. circuits. **\$165.00**

SARGENT

SCIENTIFIC LABORATORY INSTRUMENTS • APPARATUS • SUPPLIES • CHEMICALS

E. H. SARGENT & COMPANY, 4647 W. FOSTER, CHICAGO 30, ILLINOIS
DETROIT 4, MICHIGAN • DALLAS 35, TEXAS • BIRMINGHAM 4, ALA. • SPRINGFIELD, N. J.

Editorial	New Garb	829
Articles	The Villafranchian and Human Origins: <i>F. C. Howell</i>	831
	Man's bipedalism and the use of tools appeared during times of faunal change in the early Pleistocene.	
	High-Altitude Observation Techniques: <i>R. C. Staley</i>	845
	A variety of tools besides satellites are available for the study of upper-atmosphere physics.	
Science in the News	Khrushchev Sees Aspects of American Science; House Committee Questions Adequacy of Manpower Roster; Radio Frequencies for Research Studied at Geneva; Health Research Grant Practices Held Satisfactory	848
Book Reviews	A. H. Dupree's <i>Asa Gray</i> , reviewed by <i>C. Zirkle</i> ; other reviews	855
Reports	Nitrogen Fixation in Lakes: <i>R. Dugdale et al.</i>	859
	Sensory Deprivation and Visual Speed: An Analysis: <i>R. Held and B. White</i>	860
	Protection by Sulphur Compounds against the Air Pollutants Ozone and Nitrogen Dioxide: <i>E. J. Fairchild, II, S. D. Murphy, H. E. Stockinger</i>	861
	Uridine Isomer (5-Ribosyluracil) in Human Urine: <i>M. Adler and A. B. Gutman</i>	862
	Production of Large Amounts of Plant Tissue by Submerged Culture: <i>W. Tulecke and L. G. Nickell</i>	863
	Pipecolic Acid in Leaves of Strawberry Plant as Influenced by Treatments Affecting Growth: <i>L. Yatsu and D. Boynton</i>	864
Departments	Letters	826
	Forthcoming Events; New Products	866
Cover	Electron micrograph of a fractured quartz crystal, taken as part of a basic study of fracture surfaces being made at the National Bureau of Standards. Fracture surfaces are of considerable interest in determining the mode of energy dissipation once fracture has been initiated. The crystal was fractured in tension parallel to the basal planes. Replicas were made by the collodion-carbon double-replica technique, with palladium shadowing. The picture shows "steps" meeting at an angle which suggests the presence of a boundary between twin domains. Fracture markings, similar to those occurring in many materials, appear along with cleavage planes, which are of particular interest in view of the extremely poor cleavage of quartz. Such planes are less prominent in fractures propagated along other crystallographic directions. (14,700) [National Bureau of Standards]	

How the Outer Edge of the Earth's Atmosphere Can Be the Training Ground for Man's First Landing on the Moon

by Norman V. Petersen

*Chief of the Astro Systems and Research Laboratories,
Norair Division of Northrop Corporation*



One of our current studies at Norair shows that a manned capsule can be rocket-launched from the earth in a ballistic trajectory approximating an approach to the moon. A braking rocket blast fired from the capsule would push the vehicle into an earthward turn and place it in landing position above the earth's atmosphere—the same way a space ship would maneuver for a lunar landing. Such a maneuver would permit simulation of the lunar landing maneuver in near vacuum conditions as on the moon. It would utilize the blanket of air about the earth for safe recovery upon re-entry.

We base these particular studies on the use of conventional manned satellite capsules modified by the addition of a braking rocket as the lunar landing trainer and a ballistic rocket booster as the launching vehicle.

The capsule would be ejected at 50 miles altitude after traveling 100 miles from the launch site. Trajectory prior to ejection could be made to simulate either a close orbit approach to the moon, an intersecting elliptic trajectory approach or a vertical approach from a direct earth-moon trajectory. The guidance system would perform automatically during the initial approach, but would

be fitted with an "override" feature allowing the pilot to take over the controls during the braking maneuver.

For ten seconds the capsule would hover motionless above the earth's atmosphere—supported by the reverse thrust of the rocket blast. Then the pilot and capsule would re-enter the atmosphere in a low-speed free fall. After re-entry, the descent would be completed by parachute.

Ground-based stations would beam a motion picture display of moon terrain during the braking rocket descent. This, coupled with the capsule's guidance system, would give the pilot an actual impression of steering his vehicle to a moon landing.

This simulated lunar landing study is only one of our many current missile and space programs at Norair. Our range of activity allows the scientist and the engineer to work in research, design and development. He is active in the fields of close orbit, lunar and interplanetary flight regimes as well as in research in the many fields of space technology. These fields embrace astrodynamics, astronavigation, space physics, bio-astronautics, space electronics, space materials and processes, space propulsion and space structures.

Work in such challenging areas is stimulating, and the Norair environment is encouraging. For surrounding us are the finest facilities and the vast array of the entire Northrop Corporation's equipment.

The Norair scientist and the engineer enjoys deep professional satisfaction and a continuing chance to broaden his activities in space age developments. He is one of those important men who are making the news happen at Northrop.

Current papers by Northrop scientists and engineers include:

A Simulator for Lunar Landings

—Norman V. Petersen


The Influence of Launch Conditions on the Friendly Rendezvous of Astrovehicles

—Robert S. Swanson and Norman V. Petersen.


For copies of these papers and additional information about Northrop Corporation, write:

NORTHROP CORPORATION

Dept. T4-1300-32, P.O. Box 1525
Beverly Hills, California



Presenting KIMAX[®]



—new laboratory apparatus made from rugged “hard” glass

New KIMAX laboratory glassware from Kimble may offer you new opportunities for greater savings through discounts. Ask your dealer about them.

KIMAX incorporates Kimble's high standards of accuracy, design and workmanship with new durability.

26015 DISTILLING FLASK. Reinforcing bead on rim for extra strength. Side-tube seal to neck is uniform and heavy. Uniform walls, “hard” glass construction provide additional strength. In sizes from 25 ml capacity to 1000 ml.

22012 DISTILLING RECEIVER. Made

of KG-33 “hard” glass to withstand thermal shock. Calibrated to contain, retested for accuracy. Standard taper joints carefully ground. 10 ml capacity.

18130 CONDENSER. Built to withstand thermal and mechanical shock. Sealed-in inner tube has standard taper joint and drip tip at outlet. Adapter is tooled for accurate stopper fit and thick-walled for greater strength. In two sizes—300 mm and 400 mm.

24071 EXTRACTION TUBE. Made of heavy, uniform tubing for greater strength. Vapor tube protects siphon tube. Standard taper joints. Available

in three inside tube diameters—30, 38 and 50 mm.

20007 ASTM GRADUATED CYLINDER. Calibrated to contain. With round base (60 mm diameter) to fit cooling baths. Reinforcing bead at top. Uniform, heavy walls. Assured accuracy of entire scale. Durable markings. Complies with ASTM requirements. Also available in 25 ml capacity.

See your Kimble dealer for more details or write Kimble Glass Company, subsidiary of Owens-Illinois, Toledo 1, Ohio.

KIMAX is available through dealers in the United States, Canada and principal foreign cities.


KIMBLE LABORATORY GLASSWARE
AN **①** PRODUCT

OWENS-ILLINOIS
GENERAL OFFICES • TOLEDO 1, OHIO

ROYAL PRECISION

LGP-30

**Your
greatest
computer
value
on
the
market...**




inside... The world's largest-selling engineering and scientific computer, the compact, mobile LGP-30 is also the lowest-priced complete computing system you can buy or lease. It gives you the largest memory (4096 words) and capacity in its class. Internal components have been greatly reduced to insure highest operating reliability. The logic circuit and few etched circuit cards are easily removed for checkout and maintenance. **and**... The LGP-30 operates from any wall outlet, requires no site preparation. With its simplified command structure, the LGP-30 is the easiest computer to program and operate. It is delivered complete with tape typewriter for alpha-numeric input-output. An extensive library of programs and sub-routines is available. Sold and serviced coast-to-coast, in Canada and abroad by Royal McBee, it's your greatest computer value inside and **out.**



Royal Precision Corporation



Royal Precision is jointly owned by the Royal McBee and General Precision Equipment Corporations. LGP-30 sales and service are available coast-to-coast, in Canada and abroad through Royal McBee Data Processing offices. For complete information write **ROYAL MCBEE CORPORATION**, data processing division, Port Chester, New York



Chromatograph-pure fractions – Isolation of pure components, as small as 0.0002 cc, is made possible by the Beckman GC-2 Gas Chromatograph, using a new system of fraction collection. Pure, minute fractions are captured and condensed in a fluted glass tube, then transferred to microcells for infrared analysis or retained for other studies. Large samples can be introduced and fractionated with no loss of resolution, by using a unique large-column and time-controlled heated inlet. ☒ These and other accessories – such as the new Dual Column Valve, Precision Liquid Sampler, Back Purge Kit, Large-volume Gas Sampling Assemblies and Interchangeable Columns – offer new versatility and precision in gas chromatography. For details see your Beckman dealer or write for Data File 38-39-04

Beckman


Scientific and Process Instruments Division

Beckman Instruments, Inc.

2500 Fullerton Road,


Fullerton, California

PHOTO BY WILL CONNELL



- 1. NO POCKETS** to catch the product. Ball-and-socket connections let product flow straight through without hold-up.
- 2. COMPACT** ball and socket joints minimize "dead space" between columns and condensers. Entire Bantam-ware unit occupies only 2.5 sq. ft.
- 3. BETTER CLAMPING**—Outer walls of the π joints are cylindrical; permit vice-like gripping by clamps.
- 4. MOUNTING FLEXIBILITY**—Ball and socket joints prevent glassware strain while clamping.
- 5. NO-SLIP HOSE CONNECTORS** Serrated design resists slippage due to internal pressure, eliminates makeshift wired-on assemblies.
- 6. SPECIAL π JOINTS**—No product-catching bulb below grinding. Outer walls cylindrical for easy clamping. No thin spots.
- 7. CORRECT BULB POSITIONING** Adapters permit proper positioning of thermometer bulbs in relation to sidearms, thus assuring accurate temperature readings.
- 8. VERSATILE THERMOMETER JOINTS** Use π 10/18 or π 10/30 thermometers. Adapters, distilling heads, flasks, etc., are designed to take either.

ORDER this time-saving, work-saving Bantam-ware kit for fractionation, distillation, chromatographic analysis, extraction, filtration, reflux and separation. Catalog No. K-27010: \$440 complete. Write for free Catalog TG-15A.



KONTES GLASS COMPANY

First Choice For Quality Technical Glassware
Vineland, New Jersey

Midwest Distributor: Research Apparatus, Inc., Wauconda, Ill.

Letters

Comfort and Environment

Your "Dog days" editorial [*Science* 130, 131 (1959)] indicates adverse reaction to the "discomfort index" for business reasons. There are additional grounds for dispensing with the new discomfort index. Since human comfort is a rather complex function of ambient temperature, humidity, wind, and radiation load (principally solar, but also infrared as represented by wall temperature in a room), it would appear that a discomfort (or comfort) index should include the effects of all four environmental quantities. Additionally, however, the large individual differences in personal reaction to the physical environment, with both physiological and psychological factors, would appear to make determination of a rational discomfort index a virtual impossibility.

KENNETH H. JEHN

Electrical Engineering Research
Laboratory, University of Texas,
Austin

Properdin

Having read with great interest D. W. Talmage's "Immunological specificity" [*Science*, 129, 1643 (1959)], I would like to comment on the hypothesis raised, in reference to the properdin system.

No really satisfactory definition of properdin is available at the present. Although most investigators consider it a discrete entity—"a naturally immune factor"—recent work, especially that of Nelson (1), attempts to erase the distinction between the properdin and the classical antibody system. Properdin is considered a "natural antibody of broad if weak spectrum of activity." Although properdin does seem to possess certain distinct physicochemical characteristics, yet discrepancies are found in its immunological activity. For instance, there is lack of correlation of its activity as measured by phage neutralization, zymosan titration, or antimicrobial activity (2).

H. Isliker of Bern, Switzerland, indicated recently that properdin is a polymer of 7-S globulins linked reversibly by disulfide bonds (3). Thus it may be that the wide spectrum of properdin activity is due to the different affinities of component "natural globulins," and that the lack of correlation in measurements of its activity made by different methods results from variation in the ratios of immunologically different glob-

ulins incorporated into the macro-molecule.

I have always been impressed by the fact that properdin titers are consistently found to be decreased in infections and malignancies and in military personnel subjected to a massive immunization program (4)—all conditions involving a specific antibody formation. As this had never been adequately explained, we felt that something in the nature of a "competitive equilibrium" was involved—properdin polymer dissociating into its component globulins which were then "retooled" to fit the antigens flooding the organism. With Talmage's hypothesis, such a "retooling" would not be deemed necessary, with natural globulins reaggregating into specific complementary patterns. This hypothesis should certainly be easy enough to check experimentally with labeled globulins.

I certainly agree with Talmage that it would be of interest to check whether properdin could be inhibited by the mechanisms of immune tolerance. Zymosan would seem to be the logical inhibitory agent. Such an experiment would be helpful in elucidating the true nature of properdin. A finding of complete suppression would tend perhaps to strengthen the position of those who consider it a natural antibody. Also, it would provide an opportunity to ascertain properdin's role as a defense mechanism.

MICHAEL W. RYTEL

Wistar Institute,
Philadelphia, Pennsylvania


References

1. R. A. Nelson, *J. Exptl. Med.* 108, 515 (1958).
2. K. M. Cowan, *Science* 128, 778 (1958).
3. H. Isliker, *Proc. Intern. Congr. Biochem.*, 10th Congr., Vienna (1958).
4. I. Schultz and G. H. Stollerman, *Clin. Research Proc.* 5, 304 (1957).

Without commenting directly on Rytel's interesting hypothesis of the nature of properdin, I would like to clarify an apparent misunderstanding of the word *combination* as used in my article. I used the word in its abstract mathematical sense to indicate the number of different ways that independently reactive globulin molecules can be grouped. There was certainly no intention of implying a physical aggregation of natural globulin molecules. Studies with sulfur-35 amino acid incorporation in a number of laboratories have clearly shown that antibodies are formed directly from amino acids *de novo*—that is, not from preexisting globulin or other protein precursors.

DAVID W. TALMAGE

University of Colorado Medical Center,
Denver



Arc power, typified by Avco Model PG-500 Plasma Generator (above), duplicates searing heat of re-entry. Operating simplicity of Avco Model PG-030 (inset) permits wide use in the development of plasma spray techniques. Lightweight, compact—only 2¼" x 9".

NOW AVAILABLE...

USE-PROVEN PLASMA GENERATORS that duplicate the searing heat of re-entry

Avco Model PG-500 Plasma Generator, operating at one atmosphere of pressure, produces enthalpies from 250 to 12,500 Btu/lb, consumes up to 1.5 megawatts of power, uses air as the working fluid and incorporates over two years of re-entry simulation experience. PG-500, producing heat fluxes and temperatures on a *continuous* basis formerly attained on an intermittent basis in the shock tube, now provides the long running times in air necessary for materials development.

Another unit in the Avco-developed family of plasma generators, Model PG-030, consumes from 2 to 20 kilowatts of power and produces temperatures over the range from 2500°F. to 12,000°F. Model PG-030 is used for development of plasma spray techniques and spectroscopic studies. Using argon as the working fluid, it is capable of virtually unlimited operation at a power level of 15 kilowatts. Model PG-030 is especially suited to low cost

operation and features automatic starting, a high degree of controllability, quiet operation and shop-type reliability.

Write for bulletins describing Avco's products and systems in the fields of Major Research Facilities, Hypervelocity Instrumentation and Environmental Test Equipment. Write to: Products and Services Dept., Research and Advanced Development Division, Avco Corporation, 201 Lowell St., Wilmington, Mass.

Avco

Research & Advanced Development





Albert Einstein...on the chief interest of science

"It is not enough that you should understand about applied science in order that your work may increase man's blessings. Concern for man himself and his fate must always form the chief interest of all technical endeavors, concern for the great un-

solved problems of the organization of labor and the distribution of goods—in order that the creations of our mind shall be a blessing and not a curse to Mankind. Never forget this in the midst of your diagrams and equations."

—Address, California Institute of Technology, 1931


THE RAND CORPORATION, SANTA MONICA, CALIFORNIA

A nonprofit organization engaged in research on problems related to national security and the public interest

PAUL
CHAU
WALL
H. B
GEOR
MARG
THOM
DON
MINA
WILL
ALAN
PAUL
DAEL

DON
KON
EDWI

JUNE
S. D
OLIV
MUR
MOU
WORT
EA



SC
SCI
day
vanc
Com
publ
Wash
matt
is ir
Liter

En
ence
Mas
Man
ing
assu
man
cont
prep
Scie

D
addr
St.,

C
to 1
D.C
addr
and
any.

A
\$1.5
Cab

AMERICAN ASSOCIATION
FOR THE
ADVANCEMENT OF SCIENCE

Board of Directors

PAUL E. KLOPSTEG, *President*
CHAUNCEY D. LEAKE, *President Elect*
WALLACE R. BRODE, *Retiring President*
H. BENTLEY GLASS
GEORGE R. HARRISON
MARGARET MEAD
THOMAS PARK
DON K. PRICE
MINA REES
WILLIAM W. RUBEY
ALAN T. WATERMAN
PAUL A. SCHERER, *Treasurer*
DAEL WOLFE, *Executive Officer*

Editorial Board

DONALD J. HUGHES H. BURR STEINBACH
KONRAD B. KRAUSKOPF WILLIAM L. STRAUS, JR.
EDWIN M. LERNER EDWARD L. TATUM

Editorial Staff

DAEL WOLFE, *Executive Officer*
GRAHAM DUSHANE, *Editor*
JOSEPH TURNER, *Assistant Editor*
ROBERT V. ORMES, *Assistant Editor*
JUNE G. BANDY, CHARLOTTE F. CHAMBERS, SARAH
S. DEES, NANCY S. HAMILTON, WILLIAM HASKELL,
OLIVER W. HEATWOLE, YUKIE KOZAI, ELLEN E.
MURPHY, BETHSABE PEDERSEN, NANCY L. TEI-
MOURIAN, MARIA A. WOLSAK, LOIS W. WOOD-
WORTH
EARL J. SCHERAGO, *Advertising Representative*



SCIENCE, which is now combined with THE SCIENTIFIC MONTHLY, is published each Friday by the American Association for the Advancement of Science at National Publishing Company, Washington, D.C. The joint journal is published in the SCIENCE format. Entered at the Washington, D.C., Post Office as second class matter under the Act of 3 March 1879. SCIENCE is indexed in the *Reader's Guide to Periodical Literature*.

Editorial and personnel-placement correspondence should be addressed to SCIENCE, 1515 Massachusetts Ave., NW, Washington 5, D.C. Manuscripts should be typed with double spacing and submitted in duplicate. The AAAS assumes no responsibility for the safety of manuscripts or for the opinions expressed by contributors. For detailed suggestions on the preparation of manuscripts and illustrations, see *Science* 125, 16 (4 Jan. 1957).

Display-advertising correspondence should be addressed to SCIENCE, Room 740, 11 West 42 St., New York 36, N.Y.

Change of address notification should be sent to 1515 Massachusetts Ave., NW, Washington 5, D.C., 4 weeks in advance. If possible, furnish an address label from a recent issue. Give both old and new addresses, including zone numbers, if any.

Annual subscriptions: \$8.50; foreign postage, \$1.50; Canadian postage, 75¢. Single copies, 35¢. Cable address: Advancesci, Washington.

New Garb

From the time when the contents of *The Scientific Monthly* and those of *Science* were combined in January 1958, the number of pages of both text and advertising has gradually increased. The first of these increases is a consequence of deliberate policy on our part: we thought it both desirable and obligatory to expand the text of the combined magazine to approximate the total that had previously been printed in both.

The second increase, that in advertising, is attributable in part to the sharp rise in circulation that was an automatic consequence of the combination of the circulations of the two magazines: circulation of *Science* jumped from less than 40,000 to nearly 60,000. It has now passed the 62,000 mark.

The increase in size of individual copies (the 48-page issue typical of 1957 has given way to the 64-page issue typical of today) and the greatly increased press run have nearly doubled the size of our printing job. The demands of both economy and speed could be satisfied only by the use of larger and faster printing equipment. Consequently, we were reluctantly forced to shift our printing from Business Press, Lancaster, Pa., to the National Publishing Company, Washington, D.C. It was no easy or happy decision for us to terminate our cordial and effective relationship with the Business Press, a press that has performed with consistent high standards of presswork, prompt attention to our needs, and notable devotion to the job of getting *Science* out on time, despite power failures, blizzards, and other hazards.

The shift in printers gave us an opportunity to make a fresh appraisal of the magazine and to consider what changes would be desirable in the reader's interest. We decided to continue to use the same format, which we think is a good one, but to make changes in headings and type faces that would increase readability and help the reader locate material of interest. Thus the general appearance of the pages remains the same, but the type faces are different and, we hope, more readable.

The most immediately obvious change is in the cover. Formerly we used a self-cover (a cover of the same paper stock as the rest of the magazine), but as the magazine increased in weight this became progressively less satisfactory. The separate cover of heavier stock has several advantages: it will permit us to mail *Science* unwrapped and unfolded, and it will permit us to carry cover pictures of subjects of scientific interest. Some of these we will draw from articles that we are publishing, but the great majority will have to come from other sources. We invite our readers' cooperation in supplying us with suitable pictures.

The new cover has one disadvantage: it no longer carries the list of contents, a feature of the former cover that was appreciated by many subscribers. To make the best compromise possible, we have transferred the former cover material in essentially unchanged form to the third inside page. In subsequent issues it will occupy the same position.

So much for the changes. We feel somewhat like an elderly lady (after all, we were born in 1883) who has ventured to appear in a somewhat more daring gown than has been her custom. Like such a lady we hope our new garb meets with favor; unlike her we won't mind sharp and constructive criticism.—G.DuS.

Here Is A Complete Gamma Spectrometer



This transistorized spectrometer is by far the smallest and most compact available—yet it has more important and unique features than any other on the market today. Furthermore, it can be used directly with the Packard Auto-Gamma Sample Changer for completely automatic counting of test tube samples.

Write for Complete Information. Request Bulletin 400.

Packard Instrument Company, Inc.
P. O. BOX 428A • LA GRANGE, ILLINOIS

CURRENT PROBLEMS IN RESEARCH

The Villafranchian and Human Origins

Man's bipedalism and the use of tools appeared during times of faunal change in the early Pleistocene.

F. Clark Howell

Few people lack curiosity about their ancestors, their genealogy, and the ways of the world in the past. This curiosity extends to the ancestry and history of all mankind. In the last hundred years theological explanations have—especially in countries of Western cultural tradition—tended to be replaced by a more rational approach to the matter. This approach is the concern of human paleontology, and its inception is closely linked with the name of Darwin.

A century ago only a few early human skeletal remains were known, and these only from Western Europe. These represented either the approximately 35,000-year-old Cromagnon people or the (then scarcely recognized) 50,000-year-old "classic" Neanderthal folk. Much understanding of the course of human evolution has been gained since Darwin's time, and especially within the past quarter century (1). There are still some extraordinary gaps in the fossil record of the family Hominidae with respect both to specific ranges of Pleistocene time and particular geographic areas. Some such gaps, like the middle Pleistocene range in Mediterranean Africa, have become known only in the last decade, as investigations have been vigorously pursued in the field. However, fossil hominid specimens are still almost entirely unknown

from this range of time in Western Asia and in sub-Saharan Africa. The most striking hiatus, particularly since stone artifacts are far from uncommon, is the complete absence of hominid skeletal remains from the Pleistocene in the Indian subcontinent.

In spite of these deficiencies in knowledge, a good deal is known of the major evolutionary stages in hominid phylogeny for the latter half of the Pleistocene. Several distinct, largely geographically restricted lineages are recognized to have existed during the middle, and to have persisted into the early upper, Pleistocene; these probably represented paleospecies (2), although some workers consider them to have been generically distinct (3). From a portion of one such (*not* Asian) lineage, anatomically modern man (*Homo sapiens*) evolved, but the details of this transformation are still largely obscure (4). This best documented aspect of human paleontological knowledge encompasses only the later phases of man's evolution. It is well known largely because of its recency and because there are apparently more abundant traces of human occupation in datable Pleistocene contexts, especially since these can often be linked with effects of the extensive continental glaciations of the Northern Hemisphere. This record thus begins, broadly, during the time of the first of these great continental ice sheets,

variously named in Germany (5), Britain (6), Poland (7), and European Russia (8).

The main lines of hominid evolution were set during an earlier segment of the Pleistocene. The mammalian faunas of that time range, in Europe, Asia, and Africa, reveal the first appearance of new and modern genera in an otherwise often archaic assemblage. This was an extended period of fluctuating, but generally cooler and more temperate, climates compared with the late Tertiary (Pliocene), leading to mountain glaciation and changing biotopes and biotas. It was a time marked by extensive mountain building, faulting, and upwarping to the extent of several thousand meters, as seen in the Alps, Pyrenees, Caucasus, Atlas, and Himalayan ranges (9). This long interval, representing the lower Pleistocene and often termed the Villafranchian after its characteristic fauna, encompasses a time probably as long as the whole of the middle and upper Pleistocene. Half a million years is perhaps a modest estimate. The recent work by Emiliani (10), who employed oxygen isotopes in the analysis of climatic change from ocean-bottom cores, gives some promise of providing an "absolute chronology" for the Pleistocene. However, the published climatic curves appear to encompass only the latter half of the Pleistocene—that is, the three major continental glaciations and the intervening Great and Last interglacial stages, the stratigraphy of which closely parallels the core profiles.

The lower Pleistocene represents perhaps the most crucial period for future research in human paleontology. Such efforts promise results of great significance to the understanding of formative phases of hominid phylogeny and the elucidation of those distinctive associated patterns of behavior which differentiated the first hominids from other higher primate (pongid) antecedents and collaterals. I have attempted in this article to indicate a part of what is known of the Villafranchian stage from traces of hominids of this period.

The author is associate professor of anthropology, University of Chicago, Chicago, Ill.

Villafranchian: Europe and Asia

The base of the Pleistocene is best defined by three lines of evidence: tectonics, climatic deterioration, and the appearance and distribution of new forms of animal life. In the majority of stratigraphic sections, an unconformity, representing an interval of uplift and erosion, separates terminal Pliocene deposits from overlying marine or continental (Villafranchian) sediments of the basal Pleistocene. The first signs of marked cooling are evident in the appearance of north temperate or arctic forms (such as *Cyprina islandica* in

Mediterranean waters) in marine invertebrate faunas, and in vegetation changes demonstrated by palynology; or by the particular conditions of sedimentation. Continental deposits of basal Pleistocene age contain mammalian assemblages referred to as Villafranchian, from the type locality of Villafranca d'Asti in the upper Po river drainage basin (11). Such faunas are characterized by the first appearance of the modern genera *Elephas* (*Archidiskodon*), *Bos* (*Leptobos*) and *Equus* (and, in some areas, *Camelus*), but in association with a number of other typically late Tertiary species (12). The term

Villafranchian has also often been extended to apply to this basal Pleistocene interval as well as to the fauna. The termination of the interval is ill defined; it is probably best taken as the base of the Cromerian "interglacial" stage (13) or its marine equivalent in the Mediterranean basin, the Sicilian (14). This stage immediately precedes the first major continental glaciation.

In western and southwestern Europe (15), evidences of fluctuating sea levels, both transgressive and regressive, and of variably cooler climates, unlike the preceding Pliocene (16), are recognized at a number of localities. The tilted deep-water Plaisancian and brackish Astian sediments of the (later) Pliocene sea, which flooded many areas of lowland southern Europe which are continental at present, are unconformably overlain in a number of regions (in southern France and Italy) by the marine deposits of the basal Pleistocene Calabrian sea (17). The Calabrian, a transgressive sea which was subsequently regressive [compare the Po valley; also the Emilian stage of *Emilia* (18)], has as its continental equivalent, developed in unsubmerged and emergent uplands, a series of fluviolacustrine sands and gravels with a markedly cool temperate flora and a characteristic Villafranchian mammal fauna. The contrast with the subtropical vegetation and archaic mammal faunas of the Pliocene is striking and clearly delineated.

The comparable successions afforded by the North Sea basin, in East Anglia (19) and the Netherlands (20), and by the polliniferous clays and lignites of the lacustrine basin of Lefte (Bergamo, Tuscany) (21) illustrate the main pattern of climatic change during the Villafranchian in western and southern Europe (Fig. 1). The evidence would seem to indicate at least two major colder stages, the latter double, prior to the well-defined Cromerian interglacial stage. These two colder stages are separated by the still inadequately known Tiglian "interglacial" stage (22). This sequence is paralleled in the Rome region by the Acquatraversan and Cassian phases of colder climate which precede the transgressive Sicilian sea and the extensive eruptions of the Sabatino volcanoes (23). A very considerable body of evidence indicates several phases of fairly extensive mountain glaciation during this interval, very probably including the Donau (24) and Günz stages in the northern Alps. The




Fig. 1. Villafranchian stratigraphy and climatic change in the North Sea basin and the Lefte lake basin (Tuscany). The silhouettes indicate some of the characteristic species in particular faunal assemblages. The Lefte climate curve, based on palynological evidence, is expressed in terms of present altitude corresponding to vegetation; Lefte is 400 meters above sea level. [Compiled from various sources; see text citations]

first Himalayan glaciation is perhaps the Asian equivalent. However, there is a dearth of evidence of direct correlation of these subalpine stages either with continental Villafranchian deposits in the lowlands or with equivalent marine horizons.

The European Villafranchian lacks any trace of higher primates, and there are no stone implements testifying to occupation of the continent by hominids (25). This was probably a consequence of two factors: (i) Europe was not a primary center in the original hominid radiation, and (ii) the extent of the Mediterranean and Black seas (26), not only in the Pliocene but also during the Calabrian and Sicilian transgressions of the early Pleistocene, created impassable water gaps which effectively isolated Europe. The first evidence of hominid occupation of the European continent is well along in the earlier middle Pleistocene. It corresponds to the time of the Romanian regression (27) of the Mediterranean, when eustatic lowering of sea level attendant upon the first major continental glaciation evidently permitted expansion into Europe of those peoples [probably represented only by the Mauer (Heidelberg) mandible] responsible for the Abbevillian hand-axe industry (28).

In eastern Asia the continental Villafranchian is best known in northern China and in the southern foothills of the Himalayas (29). In northern China (30), the Pliocene, a period of dry subtropical-to-tropical climate with extensive lakes between stretches of desertic country, was terminated by diastrophic movements resulting in extensive erosion (Fenho erosion interval). These formerly widespread lakes were consequently displaced, and rejuvenated rivers and streams greatly enhanced their fluvial activities. In the synclinal basins of Nihowan (Hopei), Taiku, and Yüshe (Shansi), to mention only the best known, this erosion surface underlies a torrential lacustrine series of basal conglomerates overlain by sands, marls, and clays (lower Sanmenian series); the clays include a plant bed with cool dry flora (Taiku basin), and the series is capped with sands and silts which yield a characteristic Asiatic Villafranchian fauna (31). The entire series was tilted and is separated unconformably (by the Huangshui erosion interval) from the early middle Pleistocene red loams (upper Sanmenian series). No evidence of hominid fossils or of tool-making ac-

tivities have been recorded from the earlier, Villafranchian, beds. The first evidence of hominid occupation is revealed at locality 13, Choukoutien, correlative with the upper Sanmenian series on faunal grounds, which has yielded a single small chert chopping tool but no human skeletal remains.

The entire Pliocene and lower Pleistocene succession is magnificently represented in sub-Himalayan northwest India, both in the Potwar region (Punjab) to the northwest and the Siwalik Hills to the east (32, 33). The Pliocene, exposed in the middle Siwaliks series, is represented by fresh-water sandstones and shales, 6500 to 10,000 feet in thickness, deposited under tropical to subtropical climates trending toward increased aridity. The entire series was tilted during a major phase of mountain building followed by an interval of severe erosion at the end of the Pliocene. These movements, either through the development of anticlines or as a result of faulting, created a series of depressions trending northeast-southwest which were filled during the lower Pleistocene with great thicknesses of alluvial sediments (upper Siwaliks series) derived from the adjacent uplands. The earlier Tatrot zone of this series is characterized by a thick (100-foot) basal conglomerate overlain by nearly a thousand feet of coarse sandstone; interspersed silty or conglomeratic horizons contain a (still inadequately known) fauna which is characteristically Villafranchian (34, 35). The overlying Pinjor zone, some 1000 to 1500 feet of laminated silts and sands, clearly distinguished lithologically from the Tatrot zone, appears to have been deposited under warmer and more temperate conditions by sluggishly meandering streams, in contrast to the great alluvial activity that accompanied the deposition of the preceding Tatrot. A rich Villafranchian fauna is represented in the Pinjor zone (34, 36). However, neither the Tatrot nor the Pinjor provide hominid remains, nor is there evidence of stone tools testifying to hominid occupation. Such tools first appear, as the "Pre-Soan" (Punjab flake industry) chopper-chopping-tool assemblage (33, 37), in the overlying boulder conglomerate, attributed to the second Himalayan glaciation, of early middle Pleistocene age.

Discussion of the southeast Asia area of the Sunda shelf, an extension of the mainland in middle and later Pleistocene time, is less pertinent here be-

cause of the extensive Pliocene submergence. Only isolated mountain peaks were emergent along the southern (Zuider Mountains) and northern (Kendeng Hills) coasts in eastern Java, and some few uplands in the western region (38). A Villafranchian fauna is known from fresh-water sandstones and coarse conglomerates at several localities where these overlie Pliocene marine beds (39). However, skeletal remains of two distinct hominids (*Meganthropus*, *Homo modjokertensis*) first appear only in the early middle Pleistocene (40). Sundaland, including Sumatra, Java, and Borneo, and much of the present ocean floor had become largely continental by that time as a result of further uplift and marine regression consequent upon continental glaciation in the Northern Hemisphere. There is no evidence to suggest hominid occupation of the Sunda shelf during the Villafranchian stage.

The present evidence would appear to indicate that continental Eurasia was not occupied by hominids during the lower Pleistocene—that is, the Villafranchian stage. The evidence is admittedly and necessarily of a negative nature, and further field investigations, particularly in the sub-Himalayan Siwaliks, are sorely needed. The important fact is that the available evidence does not bear out the opinions of certain earlier workers (41) that Asia, or central Asia in particular, was a primary center for hominid (or higher primate) origin and dispersal. Such a conclusion is contradicted not only by our understanding of higher primate relationships, based on comparative anatomical and paleontological studies, but also by biogeographical and paleogeographical conditions. On the other hand, a variety of evidence shows that the Villafranchian stage in Africa was crucial in the earlier phases of hominid evolution.

African Villafranchian

Mediterranean Africa. Several lower Pleistocene localities with Villafranchian faunas are known in northwestern Africa (42) (Fig. 2). One of the best stratified localities is Fouarat (near Port Lyautey, Morocco), along the southern border of the Rharb plain. A Villafranchian fauna is present here in coarse sands and sandstones representing a littoral facies of a Calabrian gulf which in places filled depressions in Pliocene marine sediments (43). Hominid occu-

EASTERN ATLANTIC		WESTERN MEDITERRANEAN		
MOROCCO		ALGERIA		TUNISIA
STRATIGRAPHY		STRATIGRAPHY		STRATIGRAPHY
BASAL PLEISTOCENE	Marine conglomerates, sandstones, +90 -100 m. (Sidi Messaoud, Cap Cantin, Mazagan)	SICILIAN SEA	Marine conglomerates, sandstones +90-100m. (Sahel, Oran, Arzew, Oued Isser) (overlies tilted Bel Hachel Villafranchian)	High-level (+100m.) conglomerates and gravels, calcareo-cemented, in oueds. (no marine Sicilian)
	Mouloyian Pluvial. Cooler. +150 m. terrace erosion uplift	MARINE REGRESSION	Setifian Lake, St. Arnaud: Bel Hachel, Ain Hanech and Oranion Sahel and Algerian Sahel.	Ichkeul Lake Top not observed
	Reddened loams (Marmora) 50 m. Conglomerates (Rharr, Arbaoua) 100 m.		Traces of marine sediments +200-300 m., in Oran, Mostaganem, Chelif plain. Lacustrine depressions in Chelif and Ain Boucherit	Sandy-clays with thin sandstones (4)
	shelly-sandstones Moghrebian Transgression +300m. (warm, impoverished fauna; 50% Med. species) reddish shelly gravels and conglomerates	CALABRIAN SEA	Deep series (+100m.) of marls and lacustrine sandstones, with intercalated fossiliferous sandstones, gravels, silts, and calcareous conglomerates.	Fossiliferous sands clay lens clay horizon and clay horizon gravels Basal conglomerate. 30 cms.
PLIOCENE	Plaisancian-Astian sediments marls; sands and gravels.	PLIOCENE SEA	Plaisancian-Astian sediments marls; sands and gravels. (Chelif)	Plaisancian-Astian sediments marls; sands and gravels.

Fig. 2. Villafranchian stratigraphy in northwestern Africa. The silhouettes indicate some of the characteristic species in particular faunal assemblages. [Compiled from various sources; see text citations]

pation of the area is perhaps first recorded at a slightly later stage, probably corresponding to the regressive Emilian. Flaked pebbles, thought by some to represent primitive pebble tools, have been collected from the Arbaoua conglomerates, of basal Pleistocene age (44). These deposits, like the reddened Marmora sandy loams, mantle this region and (broadly) represent the continental equivalent of the Calabrian (Maghrebian) transgression (45). However, the artificial nature of the specimens is difficult to confirm when they are discovered in this gravel context.

The Villafranchian fauna of North Africa is more adequately known in Tunisia (Garaet Ichkeul), and particularly from old lake basins in northern Algeria (Bel Hachel; St. Arnaud). The base of Lake Ichkeul (near Ferryville) comprises southerly tilted deep-water Plaisancian and lagoonal Astian marine sediments which are overlain by fresh-water lake beds of basal Pleistocene age. The lower sands and gravels of the latter, separated from the Pliocene sediments by a thin conglomerate,

are richly fossiliferous (46); a mild temperate flora, but with substantial boreal elements, occurs in intercalated, more or less sandy, clays (47). However, the full succession at Ichkeul is still poorly known, and the faunal assemblage is incomplete, since the beds are only partially exposed during times of low water, on the northern foreshore of the present lake. At Bel Hachel (48) emergent dune sandstones, which concordantly overlie transgressive Plaisancian-Astian sediments, are eroded and filled with alluvial, weathered and reddened, conglomerates which contain terrestrial and fresh-water molluscs and a Villafranchian fauna. The more than 100-meter-high Sicilian beach rests horizontally and unconformably on the tilted and compressed series.

On the Constantine-Setifian plateau near St. Arnaud the earlier Pleistocene is exposed in deep ravines dissecting thick marls and other calcareous clayey sediments, with intercalated gravel and conglomeratic horizons, which fill old marshy or lacustrine depressions. In the Oued Boucherit (49), two horizons, separated by a meter of sterile brown

clay, contain a Villafranchian mammal fauna. The lower horizon (Ain Boucherit) is a coarse calcareous conglomerate, and the upper horizon (Ain Hanech), a cracked clay, rather sandy or with light gravels at the base. A quantity of undoubtedly primitive stone implements have been recovered here, largely from the upper horizon (50). These specimens (Fig. 3) are fashioned from naturally worn dolomitic limestone pebbles, exhibiting fresh, concave flaking scars and ranging in size from that of a tangerine to that of a good-sized orange. They are either battered over most of the surface to a multifaceted polyhedral form (boules) or are flaked along a margin unilaterally or bilaterally to produce an irregular sinuous edge characteristic of choppers or chopping tools. Unfortunately, the Ain Hanech site has been worked only briefly, since both political circumstances and the proximity of a Moslem cemetery have prevented extensive excavations.

No hominid skeletal remains have yet come to light in the formations of the North African Villafranchian. There

is unquestionable evidence of hominid occupation of this region during the terminal phases of the Villafranchian, and a site such as Ain Hanech might one day provide fossilized remains of the creatures themselves.

Sub-Saharan Africa. The central African Pleistocene was initiated by prolonged and extensive uplift, 1000 to 1500 feet at least, accompanied by downwarping in adjacent areas and by fracturing along ancient troughs. Extraordinary volcanic explosions, especially of tuffs and ashes, also took place along the eastern Rift Valley. As a consequence, the mid-Tertiary surface of erosion, whose relief was relatively gentle even during the end-Tertiary phases of valley incision, was severely deformed, and drainage systems were disrupted and even reversed (51, 52). Extraordinary depths of lacustrine deposits and subaerial, partially volcanic-derived, sediments were accumulated. These are particularly well preserved in the western (Albertine) and eastern (Gregory) rift valleys and represent good exposures of the continental Villafranchian (Fig. 4).

The Villafranchian, with an overlying middle Pleistocene series (Rawe beds), is also represented along the southern shores of a minor rift valley trending east-west, now Kavirondo Gulf, an eastern embayment of Lake Victoria (53). At Kanam, along the north and northeast slopes of the extinct volcano which forms Homa mountain, a great thickness (more than 100 feet) of lacustrine brown clays, with intercalated fine, laminated stony tuffs from intermittent volcanic explosions, have provided a good Villafranchian assemblage. A few pebble tools have also been recovered from such horizons. A small fragment of a hominid mandible is also known (54); although it was once believed to date back to the lower Pleistocene, it is now known to be of considerably more recent age.

In the Albertine rift valley an extensive lake existed in the basal Pleistocene, as indicated by the massive (at least 1500 to 2100 feet, to judge from borings) tilted and contorted beds of the Kaiso series (55). These are particularly well exposed along the margins of Lake Albert and its southern tributary, the Semliki River, as well as along the northern shores of Lake Edward (56) and in the region south of the latter adjacent to the volcanic highlands north of Lake Kivu. The beds seem to attain their maximum exposed thickness

in the southerly reaches of Lake Albert and at the adjacent mouth of the Semliki valley; thinner exposures in the upper Semliki and the northern reaches of Lake Edward may merely reflect earlier, more prolonged and intense subsidence of the rift floor in this region.

The Kaiso series is complex, and three main stages of sedimentation have been distinguished (57). Only the basal stages are generally regarded as lower Pleistocene; the upper stage is thought to be earlier middle Pleistocene. The lower stage (100 to 120 feet thick in the northern reaches of the valley) is largely silty, with some minor gravel horizons. In some localities this stage is found to overlie a basal ironstone horizon capped by unstratified sands; the ironstone seems to represent a laterite capping the down-faulted peneplain surface and provides an important datum point. This earlier stage is essentially nonfossiliferous. The middle stage (300 to 600 feet thick in the lower reaches of the valley) is predominantly clayey, with selenite evaporites and zones of gypsum. It is characterized by fine sands and sandstones and by discontinuous ironstone horizons and limonite lenses thought to represent desiccated pools. These horizons occasionally provide silicified wood and have yielded a typical, though small (13

species), Villafranchian mammal fauna. Several such ironstone "bone-beds" appear to be present, though they are restricted largely to the middle Kaiso stage (58). However, another, perhaps lower, bed is also present in the lower Semliki valley. The site of Kanyatsi (59), along the northern shore of Lake Edward just east of the Semliki outflow, has yielded traces of worked stone implements adjacent to an ancient subaerial soil horizon within the middle Kaiso stage. The specimens represent fresh flakes of quartz and quartzite, but the cores from which these were struck are apparently absent, and no pebble choppers or chopping tools have yet been found.

The central region of the Gregory rift valley has failed to provide certain evidence of lower Pleistocene formations; this is perhaps because there was extensive uplift during the later Pleistocene, or because there was relatively little deposition but considerable volcanic activity. The northern and southern reaches of the valley do afford such exposures. The tilted and step-faulted beds of the Omo series, just north and northwest of Lake Rudolph, testify to an ancient and extensive lower Pleistocene lake. A rich Villafranchian fauna occurs in extraordinary profusion in sandstone horizons intercalated in a massive succession of lacustrine vol-



Fig. 3. Pre-Chelles-Acheul implements from the site of Ain Hanech (Algeria). ($\times \frac{1}{2}$) [Courtesy C. Arambourg]

canic clayey tuffs (60). Neither hominid skeletal remains nor primitive stone implements have been recovered from the Omo series, which, since their discovery by the Bourg de Bozas expedition in 1902-1903, have only been worked during one relatively brief field season. Further investigations in this most inaccessible region are surely warranted, especially since this year I recovered Pre-Chelles-Acheul artifacts from the eroded sandstones.

In the southern reaches of this rift valley the lower Pleistocene is exposed along the tributary Eyasi trough and the adjacent region to the west of the Crater Highlands. At Olduvai Gorge (61), which extends from Lake Lagarja on the Serengeti plains some thirty-five miles to the western boundary faults of the Balbal depression, the magnificent lacustrine series of the middle Pleistocene is underlain by a flow of olivine basalt, which buries or obscures earlier horizons. Slightly to the south, in the Vogel river region on the northwest escarpment high above the Eyasi graben, are the Laetolil beds, a series of upfaulted subaerially deposited tuffs (62). These appear to be earlier than, as well as in part contemporaneous with, the Olduvai series. The upper Laetolil beds have yielded a rich Villafranchian assemblage of mammals, including some microfauna (63). The lowest horizons are also fossiliferous, but the fauna is very poorly known. Pebble tools and a hominid maxilla fragment (64), the latter believed to resemble the australopithecine (*Australopithecus*) (65), attributed to these beds may in fact be of later Pleistocene age. The basal bed (Bed I) at Olduvai Gorge is now known to contain a Villafranchian faunal assemblage comparable to that from Omo rather than of middle Pleistocene affinity. Pre-Chelles-Acheul stone implements of the Oldowan industry were first found here nearly 30 years ago. In July Dr. and Mrs. L. S. B. Leakey recovered a beautifully preserved skull of a new form of australopithecine from this bed in association with an occupation surface rich in such artifacts and with the bones of small game taken by this creature (announced at the 4th Pan-African Congress on Prehistory, Leopoldville, August 1959).

Central and eastern Africa afford some of the richest Villafranchian faunal localities in the world, coupled with an excellent Pleistocene succession. The faunas from these sites differ somewhat in composition, that from the Laetolil

beds being probably the youngest, overlapping basal Olduvai, and that from Kaiso being perhaps the oldest. The Omo fauna overlaps both Laetolil and Kaiso, and that from Kanam is probably broadly equivalent (52, 66). As yet it is impossible to determine the magnitude of the climatic change that occurred during the Villafranchian of central Africa because of the effects of tectonics in this unstable region. At present only a broad block correlation with other areas of the Old World, on the basis of faunal content, can be made. Future detailed investigations of the sediments in these basins, particularly of nonsequences and old soil horizons, coupled with palynological research, promise to throw some light on this problem.

At three such localities, Kanyatsi (Lake Edward), Kanam (Kavirondo, Kenya), and Olduvai Gorge, there is clear evidence of hominids, sometimes in the form of skeletal remains and, in all cases, in the form of deliberately fashioned stone tools. Few open habitation sites of the hominids themselves have yet come to light; however, this is largely a result of too little field work in difficult regions. There is every indication that such sites will be forthcoming in the future with concentrated work by prehistorians and Pleistocene geologists. Very complete evidence of such early hominid forms, including abundant skeletal remains, is afforded by the australopithecines of southern Africa (67).

Australopithecine Sites

The australopithecines now rank among the most numerous and best known of all Pleistocene hominids. Usually classified as a distinctive subfamily (*Australopithecinae*) (68) of the Hominidae, but quite probably representing merely a distinct genus, *Australopithecus* (69), the group contains two probably subgenerically distinct forms, (*Australopithecus*) and (*Paranthropus*). On the basis of the associated faunal assemblages, it appears that the australopithecines are all probably late Villafranchian (70); however, there is a possibility that the younger form (*Paranthropus*) may have persisted into the early middle Pleistocene (71).


Australopithecines are known from five sites at three localities (Fig. 5) in southern Africa (72). With one exception (Taungs) they occur in fossilifer-

ous breccia, composed of calcite-cemented dolomite soil, which infills former caves, formed by solution or subsidence along ancient fracture planes in dolomitic limestones of the Transvaal system. Studies of the mode of formation and the sequence of infillings of these caverns has demonstrated the significance of the degree of communication of the cavern with the outside in the accumulation of sediments (73). Travertines and intercalated bands of thin gray marly breccia, representing residual calcified material from dissolution of the dolomite, accumulate prior to the formation of any substantial opening to the surface. Subsequently, as the opening becomes progressively enlarged, surface-derived material collects in sufficient quantity to represent a state of equilibrium with outside conditions. Such breccia accumulations may serve as climatic indicators, through analysis of the sand fractions of breccia residues, minus the carbonate cement derived from roof drip, and comparison with modern dolomite soils in regions of differing rainfall in southern Africa. Very satisfactory results have been thus achieved for the Sterkfontein, Swartkrans, and Kromdraai (site A) sites; the method is not directly applicable to the Lime-works Cave site, Makapansgat, where alluviation or slope-wash, in the higher levels, are complicating factors. In general, the climate appears to have been somewhat drier in this region when (*Australopithecus*) lived and somewhat, or even considerably, wetter when (*Paranthropus*) lived than it is now. Brain (73) refers the three older Transvaal sites (Sterkfontein, Lime-works, Swartkrans) to a major dry interpluvial stage (with at least three separate peaks) and the youngest site, Kromdraai, to a succeeding, wetter pluvial stage.

The Taungs site, long since destroyed by quarrying activities, was a cave formed by solution in the capping carapace of a massive basal Pleistocene travertine banked up against a dolomite limestone cliff (Campbell Rand series) (74). The filling of the cavity was calcified sandy breccia overlain by contaminated travertine with sandy lenses from which the type australopithecine remains were most probably recovered.

The fauna associated with the australopithecines is not only varied but differs from site to site. In general it comprises other primates (both rare monkeys and abundant baboons), nu-

aus-
d but
eral it
rare
, nu-
OL. 130



merous rodents, insectivores, hyracoids, lagomorphs, numerous carnivores, including hyaenids and sabretooths, suids, an extinct sivathere, equids, and numerous antelopes. The following frequencies were obtained from counts of over 7000 bone fragments, out of a much larger number, from remnants of the gray marly breccia at Limeworks Cave (75, 76): 92 percent, antelope (293 individuals; more than two-thirds medium to small, the remainder large or very small varieties); 4 percent, other ungulates (four zebras, six chalicotheres, five rhinoceroses, one hippopotamus, 20 pigs, six giraffids); 1.6 percent, carnivores (17 hyenas, one leopard, one jackal, one wild dog, one sabretooth, and nine other small and medium species); 1.7 percent, baboons (45 individuals), rare rodents (hares and porcupines), and very rare birds and reptiles (tortoise, water turtle). Five australopithecine individuals represent only 0.26 percent of the total assemblage.

The various segments of the skeleton are very unequally represented at this site. The frequency of cranial fragments is particularly high among the nonbovid ungulates (88 percent), the carnivores (75 percent), the rodents (100 percent), and the primates, including the australopithecines (95 percent). The proportions are lower (34 percent) among the antelopes (of all sizes), and there are interesting differences in frequencies of antelope postcranial elements: cervical vertebrae (7 percent), other vertebrae (5 percent), ribs, and so on (5 percent), scapulae (9 percent), innominates (8 percent), forelimbs (37 percent), hindlimbs (20 percent), feet (6 percent) (these figures do not apply to very small antelopes, which are represented exclusively by cranial elements).

Any consideration of the diet and life habits of the australopithecines must take into account the associated fauna and frequencies of preserved skeletal segments. Hence, such figures are important and are much needed from the other sites. Several possibilities exist as to the manner in which the australopithecines and the associated fauna came to be incorporated into the breccias. The sites might have been (i) natural crevices into which animals fell; (ii) crevices into which bone accumulations were swept by natural agencies; (iii) carnivore lairs into which prey or scavenged carcasses were carried; (iv) rubbish heaps; or (v) actual occupation

sites of the australopithecines. There is no evidence at any of the sites to support (i) or (ii), although the gravelly breccia at Limeworks Cave was partly fluvial in origin; there is also evidence of stratification in the upper brownish breccia at Swartkrans, probably due to deposition in isolated pools but certainly not the consequence of stream activity. It will always be difficult to decide between (iv) and (v), but the important and still unsettled question is whether the sites were occupied and the bones were accumulated by carnivores, in particular sabretooths or hyenas, or both, or by carnivorous australopithecines.

It is necessary to bear in mind that these sites are known because of commercial lime-quarrying activities. In most cases such efforts were directed toward the basal travertines, formed largely when the caves were still solution cavities. In the case of Taungs the cave was discovered as a consequence of such mining in the massive cliff-forming travertines at Buxton, in one of which the cave happened to be situated. All the fossil mammals at the Limeworks Cave, Makapansgat, have been obtained by sorting through the extensive dump heaps left behind by the miners. Only in the case of Swartkrans (77) (in part) and some of the excavations at Sterkfontein have investigations been carried out which would permit some comprehension of the fossiliferous breccias as they existed *in situ*. However, the extreme consolidation of the fossil cave earths, a consequence of calcareous cementation, necessitates the use of explosives, so results are definitely limited with respect to details of the pattern of association and the arrangements of bones in the deposits. In the present state of knowledge it is indeed doubtful if the matter can be definitely settled until a new site can be excavated, with every attempt made not to disturb the stratigraphy and fossil associations.

Dart (75, 78) has repeatedly maintained that the extraordinary accumulations of mammalian skeletal remains in the fossiliferous breccias are a direct consequence of the predatory and carnivorous habits of australopithecines. Such remains represent in his opinion not only slaughtered prey but also scavenged carnivore kills; many of the bones were useful as tools and weapons for pounding, cleaving, scraping, stabbing, and slicing. On the basis of the preserved remains inventoried at Lime-

works Cave, specialized functions have been attributed to specific bones and portions of animal skeletons that were put to use by those "flesh-eating, skull-cracking and bone-breaking, cave-dwelling apes."

There are in fact two distinct issues involved here. The question as to whether the bones were employed as implements and weapons by australopithecines presupposes that these creatures were carnivorous and were therefore responsible for the fossiliferous accumulations. The use of these bones is extremely difficult to verify, since none of the sites yield any trace of specimens which have been deliberately worked or shaped. There is no doubt that the jaws and teeth, horns, and shattered or damaged limb bones which Dart attributes to an "osteodontokeratic culture" might be employed in the fashions he has so exhaustively and imaginatively outlined. However, as in the case of the so-called "bone and antler industry" from the Choukoutien locality in northern China, attributed by Breuil (79) to the middle Pleistocene hominid found there, this is extremely difficult to confirm scientifically, even though both claims may prove entirely valid.

The question of the carnivorous habits of australopithecines is a separate matter and one which should be resolved from existing evidence. The parts and proportions of the animal skeletons preserved do coincide closely with remains at carnivore kill sites in the open, even after the usual scavengers have been at work (80). Moreover, although contrary claims have been made (81), both brown and spotted hyenas may eat and accumulate bones in and about their lairs, at least at times (82). Nonetheless, this possibility does not account for the enormous concentration of bones at the sites. Also, there are discrepancies between the proportion of cranial and postcranial elements of the antelopes compared with the other ungulates, the carnivores, and the primates.

Two main points are important in connection with the dietary habits of these creatures: (i) the evidence from Taungs, Sterkfontein, and Limeworks Cave of baboon skulls bearing evident signs of localized depressed or radiating fractures, smashed-in walls or tops of the cranial vault, openings in the vault or base, and twisted facial skeletons all testify to predatory activities which are those of a hominid rather than any

hyaenid or felid carnivore; and (ii) the substantial quantity of various antelope and other long bones which are not only broken and smashed but also split longitudinally, and which usually fail to reveal any traces of carnivore gnawing, is further testimony to hominid habits.

Such evidence has been convincing not only to me, but also to other workers (83) who have examined the specimens in question. Moreover, it seems very likely, on the basis of the Taung evidence, that eggs, crabs, turtles, birds, rodents, and smaller antelopes were a not insubstantial part of australopithecine diet. The former such items are easily collected, and it is not particularly difficult to kill members of the other species of smaller mammals. The australopithecines were very probably carnivorous predators as well as scavengers of the kills of other carnivores (of

which there was then an abundance of forms long since extinct). The marked disproportion between the bovid and nonbovid ungulates and the relatively few carnivores in the Limeworks Cave inventory may merely be a reflection of this latter fact and of the limited hunting capabilities of such creatures. Such a conclusion does not preclude the possibility that carnivores also, at least periodically, occupied such sites and contributed to the bone accumulations. This can hardly be denied until careful excavations have been carried out which prove the situation to have been otherwise.

Until recently none of the australopithecine sites were known to contain artifacts. Consequently, many workers asserted that such primitive creatures, although admittedly hominids, were incapable of making, and perhaps even of using, tools. Quite possibly this lack

of stone implements has also convinced some workers that bone, horn, and teeth of other animals were used by australopithecines as weapons and implements. A few split and flaked dolomite pebbles from the calcified stony and sandy fluviatile horizon which overlies the pink and gray breccias at Limeworks Cave (84) suggest but do not afford conclusive proof of tool-making activities. However, there is no doubt about the validity of the implements (Fig. 6), referred to as a pebble-tool (Pre-Chelles-Acheul) industry, recovered recently from the Sterkfontein locality (85). The specimens derive from a reddish-brown breccia at the extension site first thought to be broadly contemporary with the basal pink australopithecine-bearing breccia of the type site. Robinson's (86) more recent investigations indicate, however, that these breccias are separated unconform-




Fig. 5. Australopithecine localities in southern Africa. [Base map redrawn from the geological map of southern Africa in A. L. du Toit, *The Geology of South Africa*, ed. 3, 1954; insets adapted from G. B. Barbour, H. B. S. Cooke, J. T. Robinson, and F. E. Peabody]

ably, as a consequence of subsidence, and that the latter pink breccia underlies the reddish-brown breccia of the extension site. The latter contains the foreign and worked stones as well as some fauna (including *Equus*, absent in the type site basal pink breccia) and some remains (isolated permanent teeth, a juvenile maxilla fragment with several teeth) referred to *Australopithecus*.

The specimens recovered from Sterkfontein include pebble- and core-choppers (8), a chopper-hammerstone (1), and rough retouched end-struck flakes (2). Quartz, quartzite, chert, and diabase pebbles foreign to the deposit were also found; about half (24) of these were plain, and the other half (23) exhibited evidence of fracture from use. The small flakes struck from the choppers and cores are missing and indicate that the specimens were collected and worked elsewhere, near streams where raw material was available, prior to being carried to the site for use. Some of the specimens show extensive battering rather than careful flake removal, suggesting either hard use or, according to J. D. Clark (87), poor workmanship. The implements are fresh and unweathered and cannot have been washed in from the outside, since the breccias must have accumulated under an overhanging roof, and since the breccias fail to reveal such conditions of deposition. The artifacts seem to be concentrated at the western end of the site near the original entrance to the cave and were undoubtedly left behind by the hominids who occupied this end of the cavern.

The recovery of pebble tools in association with australopithecines is a momentous discovery. There now seems little doubt that these primitive creatures were already capable of using and manufacturing implements of stone and, presumably, of other nonpreserved materials as well. The extraordinary concentration of other mammalian bones would indicate that these creatures were capable of killing the moderate- and smaller-sized species; probably they also scavenged carnivore kills. There is no indication that they had the ability, the equipment, or the organization necessary for killing very large mammals, in contrast to middle Pleistocene peoples. Such carnivorous habits would have required some sharpened stone implements, such as flakes and chopping tools, for cutting open the hide of kills to obtain meat. There is no trace of the use of fire at this early time.

Clark (87) has recently suggested

that the availability of water in the cave systems may have been an important factor in their having been occupied. Except for small and seasonally dry streams, cave and fissure systems, and springs related to them, provide the best source of water in such limestone country. Such sources of water would have attracted game and australopithecines alike and would have provided ideal conditions for the latter to prey on antelopes, pigs, baboons, and other animals which came to drink there. This also readily accounts for the profusion of animal bones accumulated in the cave and for the presence of implements necessary to butcher the slaughtered game.

Australopithecine Morphology

Hominids and pongids (apes) are generally regarded as closely related higher primate groups (hominoids). There are obvious and significant morphological and behavioral differences between the living representatives of the two families. Acceptance of the fact of evolution and the reality of such close affinity indicates, however, that such divergences were fewer and less sharply delineated in the remote past. The primary adaptation of the hominid radiation required transformation of the locomotor skeleton to permit fully upright posture and an efficient bipedal gait (88). This mode of terrestrial locomotion contrasts markedly with the arboreal or terrestrial quadrupedalism of the lower catarrhine monkeys or with the arboreal brachiation of the pongids, coupled, in the larger pongine (anthropoid ape) species, with semipronograde quadrupedalism. This basic locomotor adaptation of the hominids was doubtless preadaptive for subsequent evolutionary changes which affected the skull—reduction of the facial skeleton and extraordinary enlargement of the cerebral hemispheres and cranial vault. The latter has been linked with enhanced cultural capacities, although the manifold interrelationships between structure and function, and their behavioral significance, are still very largely obscure.

The pelvis of four australopithecine individuals are now known. Three represent (*Australopithecus*) and were found at Limeworks Cave (89) and Sterkfontein (90) (at the latter site, much of an associated vertebral column was also found); (*Paranthropus*) is represented by one incomplete innomi-

nate bone from Swartkrans (91). These fortunate discoveries demonstrate conclusively that the pelvic structure of these creatures was that characteristic of primitive bipedal hominids. They also greatly add to our understanding of the hominid locomotor transformation, which involved a complex of interrelated structural modifications. These constituted a basic reorientation of the pelvis in relation to the trunk, interrelated changes which permitted an erect trunk and full extension of the lower limbs in stable upright posture (and in the female, maintenance of the bony birth canal).

Such changes involve (i) expansion of the iliac blade, especially the auricular area, coupled with sacral rotation and accentuated lumbar lordosis; (ii) shortening and anterior rotation or "twisting" of the ilium, with attendant development of a sigmoid curvature of the iliac crest; (iii) thickening of the outer bony table above the acetabulum (to aid in balance and weight support); (iv) development of an iliac cristall tubercle, in line with the strengthened supra-acetabular region (related to differentiation and expansion of the ilio-tibial tract as an aid in stabilization of the hip and knee joints in standing and walking erect); (v) enlargement and approximation to the acetabulum of the anterior inferior iliac spine (related to the size of the straight head of the *rectus femoris* muscle, as part of the general enlargement of the *quadriceps femoris* muscle group in bipeds); (vi) shortening of the ischium and altered form of the ischial tuberosity (the full significance of this is obscure, but it is apparently related to the position of the extensor, or hamstring muscle, lever arm); (vii) enlargement and displacement of the *gluteus maximus* muscle as a powerful extensor (rather than a lateral rotator as in apes and monkeys); and (viii) altered function of *gluteus medius* (and *minimus*) muscles as abductors (to maintain lateral stability in walking erect).

There are accompanying interrelated modifications in the proximal head of the femur. These include: (i) enlargement of the femoral head; (ii) development of the lesser trochanter (related to an altered disposition of the *psaos major* muscle); (iii) development of the anterior segment of the greater trochanter and the intertrochanteric line (related to the attachment of the Y-shaped ligament of Bigelow and the joint capsule); (iv) development of the *linea aspera*; (v) shift of *gluteus maxi-*

mus to a posterior rather than a lateral insertion, as in apes, in the place of *adductor minimus* and expansion of *vastus intermedius*; and (vi) notable reduction of the *quadratus femoris* muscle. Other modifications of the distal end of the femur, including the obliquity of the shaft, the marked depth of the patellar surface, the configuration of the intercondylar notch, and the enlargement of the lateral condyle, were apparently associated in large part with enhanced stability of the knee joint in orthograde progression.

The basic morphological pattern characteristic of hominid bipedalism is apparent in the australopithecine lower limb skeleton so far as it is known (92). There are a number of minor differences—for example, in the form of the ischial region—from the pelvic morphology of *Homo sapiens*, a not unexpected finding in a primitive lower Pleistocene hominid. The australopithecine lower leg and foot, except for a talus of (*Paranthropus*), is still largely unknown.

The morphological pattern of the australopithecine dentition is also hominid rather than pongid (93). In the deciduous dentition this is evident in (i) the evenly curved dental arch, lacking diastemata; (ii) the small milk incisors; (iii) the small, nonprojecting spatulate milk canines; (iv) the quadricuspid upper first milk molar; (v) the nonsectorial, quinticuspid lower first milk molar with well-developed anterior fovea and cusps of approximately equal height. In the permanent dentition this is evident in (i) the evenly curved (parabolic) dental arch; (ii) the small incisors; (iii) the small, nonprojecting spatulate canines, lacking a talonid and with the internal cingulum forming a basal tubercle; (iv) the double-rooted upper first premolar; (v) the nonsectorial, bicuspid lower first premolar; and (vi) the replacement sequence, in which both the permanent canine and medial incisor tend to erupt relatively early.

The related structure of the facial skeleton is also primitively hominid, and this is paralleled in a number of structural details of the cranial base and the occiput. The brain, in proportion to body size, in certain aspects of its form and proportions, and in its tendency toward delayed maturation, approaches a primitive hominid rather than a pongid condition.

There are consistent morphological distinctions between the earlier form (*Australopithecus*) and the younger

form (*Paranthropus*) (94). These are evident not only in the deciduous and permanent dentitions and the facial skeleton but also in the structure of the cranial base and vault, as well as in the known portions of the postcranial skeleton. (*Australopithecus*) was a small, gracile bipedal creature, weighing certainly no more than 75 to 85 pounds in the larger males. (*Paranthropus*), on the other hand, was a far more

robust and massive creature of probably half again that body weight. Such, probably subgeneric, differences indicate a pronounced bifurcation within a primary australopithecine radiation, at least in the basal Villafranchian and possibly even in the later Pliocene. Unfortunately the general absence of fossiliferous Pliocene horizons in sub-Saharan Africa has thwarted investigation of the earlier evolutionary phases




Fig. 6. Pre-Chelles-Acheul implements from the australopithecine-bearing breccias at Sterkfontein (Transvaal). [After J. T. Robinson and R. J. Mason, 85]

At Swartkrans another distinct hominid, designated *Telanthropus*, is found in direct association with the australopithecine (*Paranthropus*) (95). This form, which is still only very inadequately known, differs markedly from (*Paranthropus*) in dental and mandibular morphology and in certain features of the maxilla (in particular in the structure of the nasal floor). In all such characteristics, some of which reveal some resemblances to the earlier australopithecine (*Australopithecus*), this hominid is further evolved morphologically than any known australopithecine and approaches therefore the phylogenetic status of certain earlier middle Pleistocene forms attributed to the genus *Homo*. The full implications of this conclusion cannot be properly evaluated until additional, more complete specimens have been discovered.

It is significant that the African Villafranchian hominids differ considerably in morphology from later Tertiary hominoids of eastern Africa and Eurasia. Four hominoid genera are recognized from the earlier Miocene of eastern Africa, three and possibly four genera (96) are recognized from the middle and late Miocene and the lower Pliocene of peri-Alpine Europe, and four genera are recognized from the (upper) lower and middle Siwaliks of Asia (Fig. 7).

In both Africa and Europe primitive hylobatids (gibbons) are already evolved in the lower half of the Miocene. Most of the basic cranial and dental morphology of the group is already established in these forms; in fact the dentition is already basically hominoid in the Oligocene form *Pro-*

pliopithecus. However, in some features of postcranial structure and in limb proportions these forms differ significantly from their living representatives (97). Similarly, the basic dryopithecine (pongid) dental characteristics are manifest in the earlier Miocene hominoids of eastern Africa, although cranial and facial morphology is distinct from that of evidently specialized living varieties of the group (98). Moreover, the fundamental locomotor pattern of the recent large-bodied brachiator is evident in the morphology of shoulder and elbow joints, whereas the skeleton of the hand is distinctly primitive cercopithecoid or monkey-like (99). The later Miocene and Pliocene dryopithecine hominoids of Europe and their Pliocene counterparts of Asia, known nearly exclusively from jaws and teeth, are typically pongid in mandibular and dental structure and therein do not differ fundamentally from the living gibbons and great apes.

Oreopithecus, a form known for over three quarters of a century which has received much publicity and been much discussed recently, is a notable exception to this statement. This primate occurs in the Pontian (lower Pliocene) lignites of Tuscany (100). It is surely significant that *Oreopithecus* reveals features of dental morphology which are not typically pongid, although there can be no question that it is fully hominoid rather than primitive cercopithecoid (cercopithecoid) (101). A majority of workers tend to agree that the other hominoids of the later Tertiary are typical pongids, with all the attendant dental specializations which would effectively exclude such creatures from hominid ancestry. Hürzeler's (102) painstaking reexamination of all the *Oreopithecus* material proves, however, that all later Tertiary hominoids were not typically pongid (103). Hence, the long-standing argument over the "cercopithecoid" (Old World monkey) or "pongid" (anthropoid ape) origin of hominids can be clarified, since many of the more "generalized" hominid features which some workers have regarded as indicative of lower catarrhine affinity are present in either primitive or nonpongid hominoids. Consequently, an oreopithecine hominoid group might well provide the ancestral stage from which orthograde bipedal hominoids were subsequently to evolve. It is still premature to assert the correctness of such a hypothesis to the exclusion of all others. The recent and truly important discovery of a nearly entire skeleton of




Fig. 7. Temporal distribution of hominoids in the Neogene of Africa, Asia, and Europe.

Oreopithecus from the Grosetto lignites should provide most of the evidence necessary to resolve this problem.

It should be recalled that the whole of the African Pliocene, a span of over ten million years, is still either almost or entirely unknown (104). Late Tertiary hominoids surely occupied the more central area of the continent, although fossiliferous deposits of this time range are still unknown. Until some evidences of hominoid varieties are forthcoming from the upper Neogene of sub-Saharan Africa, any hypotheses of hominid origins will lack support. There is widespread evidence of extensive sub-Saharan desiccation, between 1°N and 20°S, from the late Miocene throughout the Pliocene. This was a period of desertification during which great distributions of fine, light, unstratified aeolian sands of the (upper) Kalahari system occurred (105). Conditions appear to have been such that, except for rare, and probably small, basins of sedimentation (proto-rift valleys), which are usually obscured by volcanic lavas and other deposits, the preservation of mammalian fossils was literally precluded.

This was certainly a crucial time for mammalian evolution in general as well as for higher primate evolution in particular. Thus, the origins of the incredibly rich eastern-central African grasslands fauna is literally unknown; yet this long interval was probably vital for such a radiation, since the Villafranchian antelopes appear largely referable to existing genera. For the pongid hominoids, whose ecological requirements and habits of locomotion were apparently becoming those of forest-dwelling, vegetarian (or frugivorous) brachiators, such desiccation had profound effects on distribution, which are fully evident today in the sparse and restricted habitats of the African apes. For those hominoids that were pre-adapted toward terrestrial bipedalism, it was a period of trial in a new and exploitable environment. The Villafranchian stage, as well as the later Pleistocene phases of hominid history, testify to the achievement of this primary radiation.

References and Notes

1. The best single, well-illustrated reference work to the hominid (and primate) fossil record is the recent work by J. Piveteau, *Traité de Paléontologie, VII: Primates; Paléontologie humaine* (Masson, Paris, 1957).
2. F. C. Howell, *European and Northwest African Middle Pleistocene Hominids*, in press.
3. W. E. Le Gros Clark, *The Fossil Evidence for Human Evolution* (Univ. of Chicago Press, Chicago, 1955).
4. F. C. Howell, *Am. J. Phys. Anthropol.* 9, 379 (1951); *Quart. Rev. Biol.* 32, 330 (1957); in, *Hundert Jahre Neanderthaler 1856-1956* (Kemink en Zoon, Utrecht, 1958), p. 185.
5. P. Woldstedt, *Norddeutschland und angrenzende Gebiete im Eiszeitalter* (Koehler, Stuttgart, 1950).
6. D. F. W. Baden-Powell, *Geol. Mag.* 85, 279 (1948); W. B. R. King, *Quart. J. Geol. Soc. London* 111, 187 (1955); R. G. West, *Quaternaria* 2, 45 (1955); —, *J. Glaciol.* 3, 211 (1958).
7. W. Szafer, *Ann. soc. géol. Pologne* 22, 1 (1953).
8. H. Spreitzer, *Quartär* 3, 1 (1941); V. Gromov, *Am. J. Sci.* 243, 492 (1945).
9. See the summary list of uplifted highlands in R. F. Flint, *Glacial and Pleistocene Geology* (Wiley, New York, 1957).
10. C. Emiliani, *Science* 123, 924 (1956); *J. Geol.* 63, 538 (1955); *ibid.* 66, 264 (1958).
11. L. Pareto, *Bull. soc. géol. France* 22, 210 (1865).
12. E. Haug, *Traité de Géologie* (Colin, Paris, 1911); A. T. Hopwood, *Proc. Geologists' Assoc. Engl.* 46, 46 (1935); — *ibid.* 51, 79 (1940). Although several mammals are included as new arrivals in the Villafranchian, all are not necessary to prove a lower Pleistocene age; thus, *Equus* never reached Java and *Leptobos* is not present in sub-Saharan Africa.
13. See C. Reid, "The Geology of the Country around Cromer," *Mem. Geol. Survey G. Brit. Engl. Wales* (1882).
14. M. Gignoux, *Compt. rend.* 147, 1497 (1908); A. C. Blanc, *Geol. Meere Binnengewässer* 5, 137 (1942).
15. In France: Sâone basin (Bresse-Dombes region), Puy-en-Velay (Haute-Loire), Allier valley (Perrier, Mont Coupet, Senèze localities), and Rhone valley (St. Vallier, Drôme); in Italy: Po valley (Piemonte, Emilia), upper Arno valley (Tuscany), Monte Mario (Rome), Puglia and Calabria, and Sicily. For an excellent general review of the evidence in southern Europe see H. L. Movius, Jr., *J. Geol.* 57, 380 (1949); also, various papers in *Rept. Intern. Geol. Congr. London, 18th Congr.*, 1948 (1950), part 9, sect. H, and A. C. Blanc et al., *Compt. rend. congr. géol. intern.*, 19^e Congr., Algiers, 1952 (1954), fasc. 15, p. 215.
16. For useful summaries of European Pliocene faunas see G. Depape, *Ann. soc. sci. Bruxelles* 48, 39 (1928); also, W. Szafer, *Inst. Geologiczny, Prace* (Warsaw, Wydawnictwa Geol.) 11 (1954).
17. M. Gignoux, *Ann. géogr.* 18, 141 (1909); *Ann. univ. Lyon* 1, No. 36 (1913); *Bull. soc. géol. France* 14, 324 (1915); *Ann. Univ. Grenoble* 28, 9 (1916).
18. R. Ruggieri and R. Sellì, *Rept. Intern. Geol. Congr. London, 18th Congr.*, 1948 (1950), part 4, sect. H, p. 85.
19. C. Reid, "The Pliocene Deposits of Britain," *Mem. Geol. Survey G. Brit. Engl. Wales* (1890); F. W. Harmer, *Proc. Geologists' Assoc. (Engl.)* 17, 416 (1902); P. G. H. Boswell, *ibid.* 63, 301 (1952); C. P. Chatwin, *East Anglia and Adjoining Areas* (Geological Survey and Museum, London, 1954); for general discussion, see W. B. R. King, *Quart. J. Geol. Soc. London* 111, 187 (1955).
20. I. M. van der Vlerk, *Proc. Koninkl. Ned. Akad. Wetenschap.* B56, 34 (1953); — and F. Florschütz, *Verhandel. Koninkl. Ned. Akad. Wetenschap. Afdel. Natuurk.* 2, 1 (1953); R. Lagaij, *Mededel. Geol. Sticht., Ser. C, No. 5* (1952); for paleogeographic maps, see J. H. van Voorthuysen, *Geol. en Mijnbouw* 19, 263 (1957); for general summary, see A. J. Pannekoek et al., *Geological History of the Netherlands* (Government Printing Office, s'Gravenhage, 1956).
21. S. Venzo and F. Lona, *Atti soc. ital. sci. nat. e museo civico storia nat. Milano* 89, 43 (1950); S. Venzo, *Geol. Bavarica* 19, 74 (1953); *Actes congr. assoc. intern. étude quatern.*, 4^e Congr., Rome-Pise, 1953 (1956), vol. 1, p. 65.
22. W. H. Zagwijn, *Geol. en Mijnbouw* 19, 233 (1957); A. Schreuder, *Arch. néerl. zool.* 7, 153 (1945).
23. A. C. Blanc, *Quaternaria* 4, 95 (1957).
24. E. Eberl, *Die Eiszeitenfolge im nördlichen Alpenvorlande* (Filsen, Augsburg, 1930); I. Schafer, *Geol. Bavarica* No. 19 (1953), p. 13. Supposedly artificially flaked specimens have been reported from the East Anglian "Crags" [see especially J. R. Moir, *The Antiquity of Man in East Anglia* (Cambridge Univ. Press, Cambridge, 1927)]; however, most workers are now convinced that these are natural specimens. See A. S. Barnes, *Anthropol.* 48, 217 (1938); *Am. Anthropol.* 41, 99 (1939); S. H. Warren, *Southeast Nat. Antiq.* 53, 48 (1948).
26. M. Pfannenstiel, *Geol. Rundschau* 34, 341 (1944); *Bonner Geogr. Abhandl.* 6, 1 (1950); S. Erine, *Rev. Geogr. Inst. Univ. Istanbul* 1, 85 (1954).
27. A. C. Blanc, *Geol. Meere Binnengewässer* 5, 137 (1942).
28. It has been maintained that the Abbevillian, at least in the Somme valley where it is best known, is of "first interglacial" (=Günz-Mindel=Cromerian) age. See H. Breuil, *Proc. Prehist. Soc.* 5, 33 (1939); — and L. Koslowski, *Anthropol.* 41, 449 (1931); — *ibid.* 42, 27, 291 (1932); — *ibid.* 44, 249 (1934). However, a variety of both geological and paleontological evidence indicates that assignment of a younger age is fully warranted, and that this was probably an ameliorative phase within or toward the end of the first major continental glaciation.
29. See the excellent general summary by H. L. Movius, Jr., *Papers Peabody Museum Am. Archaeol. Ethnol. Harvard Univ.* 19, 1 (1944); also, H. de Terra, *Proc. Am. Phil. Soc.* 77, 289 (1937).
30. P. Teilhard de Chardin, *Bull. Geol. Soc. China* 16, 195 (1937); *ibid.* 17, 169 (1937); *Compt. rend. soc. géol. France*, No. 17 (1938), p. 325; *Inst. Géol.-Biol., Pekin*, No. 7 (1941); — and C. C. Young, *Bull. Geol. Soc. China* 12, 207 (1933); G. B. Barbour, *ibid.* 10, 71 (1931); H. de Terra, *Inst. Géol.-Biol., Pekin*, No. 6 (1941).
31. See faunal list in P. Teilhard de Chardin and P. Leroy, *Inst. Géol.-Biol., Pekin*, No. 8 (1942).
32. H. de Terra, *Proc. Am. Phil. Soc.* 77, 289 (1937); — and P. Teilhard de Chardin, *ibid.* 76, 791 (1936); G. E. Pilgrim, *Records Geol. Survey India* 40, 185 (1910); W. D. Gill, *Quart. J. Geol. Soc. London* 107, 375 (1952).
33. H. de Terra and T. T. Paterson, *Carnegie Inst. Wash. Publ.* No. 493 (1939), p. 1.
34. E. H. Colbert, *Trans. Am. Phil. Soc.* 26, 1 (1935).
35. The interesting and little known pongid *Ramapithecus* is the last Siwalik primate to occur in the Tattot zone.
36. D. A. Hooijer and E. H. Colbert, *Am. J. Sci.* 249, 533 (1951); also, D. A. Hooijer, *Proc. Koninkl. Ned. Akad. Wetenschap.* B55, 436 (1952).
37. H. L. Movius, *Trans. Am. Phil. Soc.* 38, 329 (1948); *Cahiers hist. mond.* 2, 257, 520 (1955).
38. H. de Terra, *Trans. Am. Phil. Soc.* 32, 437 (1943); R. W. van Bemmelen, *The Geology of Indonesia* (Nijhoff, The Hague, 1949).
39. G. H. R. von Koenigswald, *Proc. Koninkl. Ned. Akad. Wetenschap.* B38, 188 (1935); *Quartär* 2, 28 (1939).
40. D. A. Hooijer, *Am. J. Phys. Anthropol.* 9, 265 (1951); *Proc. Koninkl. Ned. Akad. Wetenschap.* B55, 436 (1952); *ibid.* B60, 1 (1957); *Quaternaria* 3, 5 (1956). A contrary opinion is held by G. H. R. von Koenigswald, *Proc. Koninkl. Ned. Akad. Wetenschap.* B59, 204 (1956); *Actes congr. assoc. intern. étude quatern.*, 4^e Congr., Rome-Pise, 1953 (1956), vol. 1, p. 5.
41. D. Black, *Bull. Geol. Soc. China* 4, 133 (1925); W. D. Matthew, *Climate and Evolution* (N.Y. Acad. Sci., New York, ed. 2, 1939).
42. C. Arambourg, *Bull. soc. géol. France* 19, 195 (1949); "La paléontologie des vertébrés en Afrique du Nord française," *Congr. géol. intern.*, 19^e Congr., Algiers, 1952, *Monogr. Arambourg* distinguishes lower (Ichkeul, Bouchet, Fouaret) and upper (Bel Hachel, Ain Hanach) Villafranchian faunas; the former is more archaic and includes *El. africanus*, *Anancus*, *Stylohyponion*, and *Libytherium*, whereas the latter includes predominantly *El. meridionalis* and *Equus*, with *Libytherium* and, rarely, *Stylohyponion*.
43. G. Choubert et al., *Notes et mém. serv. géol. Maroc*, No. 71 (1948), p. 15; G.

- Choubert, *Notes et mém. serv. géol. Maroc* No. 76 (1950), p. 13.
44. P. Biberson et al. *Compt. rend.* **245**, 938 (1957).
 45. G. Choubert and R. Ambroggi, *Notes et mém. serv. géol. Maroc*, No. 117 (1953), p. 3; G. Choubert et al. *Compt. rend.* **243**, 504 (1956); G. Choubert, *ibid.* **245**, 1066 (1957); *Actes congr. assoc. intern. études quatern.*, 5^e Congr. Madrid-Barcelona, 1957 (in press).
 46. C. Arambourg and M. Arnould, *Bull. soc. nat. Tunisie* **2**, 149 (1949).
 47. C. Arambourg et al., *Compt. rend.* **234**, 128 (1952); *Arch. Mus. nat. hist. nat. Paris*, sér. 7 (1953), vol. 2, p. 1.
 48. M. Dalloni, *Bull. soc. géogr. archéol. prov. Oran* **61**, 1 (1940); *Actes congr. assoc. intern. études quatern.*, 4^e Congr., Rome-Pise, 1953 (1956), vol. 1, p. 19; *Bull. soc. hist. nat. Afrique nord* **45**, 134 (1954).
 49. C. Arambourg, *Bull. soc. géol. France* **19**, 195 (1949); *Compt. rend.* **229**, 66 (1949).
 50. —, *Compt. rend. soc. géol. France*, No. 7 (1949), p. 120; *Bull. soc. préhist. franç.* **47**, 348 (1950); *Compt. rend.* **236**, 2419 (1953); — and L. Balout, *Bull. soc. hist. nat. Afrique nord* **43**, 152 (1952). Crude bifaces, referable to a basal Chellean industrial stage, were discovered in 1952, derived from a higher level at the Ain Hanesh site.
 51. F. Dixey, *Quart. J. Geol. Soc. London* **102**, 339 (1946); *The East African Rift Valley, Colonial Geology and Mineral Resources Suppl. Ser.* (London, 1956). See especially the recent excellent critical appraisal by H. B. S. Cooke (52, p. 60).
 52. H. B. S. Cooke, *Annex. Trans. Geol. Soc. S. Africa* (1958).
 53. P. E. Kent, *Geol. Mag.* **79**, 117 (1942); E. P. Saggerson, *Geology of the Kisumu District* (Geological Survey, Kenya, Nairobi, 1952).
 54. L. S. B. Leakey, *The Stone Age Races of Kenya* (Oxford Univ. Press, London, 1935); compare M. F. Ashley Montagu, *Am. Anthropol.* **59**, 335 (1957).
 55. This series is named after the village of Kaiso, the type locality, on the eastern shore of Lake Albert. See, E. J. Wayland et al., *Occasional Papers Geol. Survey Uganda* **2**, 1 (1926); also, K. A. Davies, *Geol. Mag.* **88**, 377 (1951); N. Harris, J. W. Pallister, J. M. Brown, *Geol. Survey Uganda Mem.* No. 9 (1956).
 56. V. E. Fuchs, *Geol. Mag.* **71**, 97 (1934).
 57. J. Lepersonne, *Ann. soc. géol. Belge* **72**, 1 (1949).
 58. The particular horizon of the type Kaiso mammal fauna is not clear; possibly it corresponds to the middle or even the upper stage, but this problem may be settled by careful study and comparison of the molluscan faunas within the several stages of the Kaiso series. W. W. Bishop (Geological Survey of Uganda, Entebbe) is now carrying out systematic investigations in the Western Rift, both at Kaiso and in the Lake George-Lake Edward region (see "Proc. Assoc. African Geol. Services, Leopoldville, Belgian Congo, Meeting, July 1958," in press).
 59. J. de Heinzelin de Braucourt, "Le fossé tectonique sous le parallèle d'Ishango," *Publ. Inst. Natl. Parcs Nat., Bruxelles* (1950); also, *Proc. Intern. Sci. Sol.*, 5^e Congr. Leopoldville, 1954 (1954), vol. 4, p. 435.
 60. C. Arambourg, *Mission scientifique de l'Omo, 1932-1933*, I (Museum Natl. Hist. Nat., Paris, 1935; 1943), vols. 1, 3.
 61. L. S. B. Leakey, *Olduvai Gorge* (Cambridge Univ. Press, Cambridge, 1951); *Nature* **181**, 1099 (1958).
 62. P. E. Kent, *Geol. Mag.* **78**, 173 (1941); also, L. Kohl-Larsen, *Auf den Spuren des Vormenschen* (Strecker and Schroeder, Stuttgart, 1943).
 63. W. O. Dietrich, *Palaeontographica* **A94**, 43 (1942); *ibid.* **A99**, 1 (1950).
 64. A. Remane, *Z. Morphol. Anthropol.* **42**, 311 (1951); *Am. J. Phys. Anthropol.* **12**, 123 (1954).
 65. J. T. Robinson, *ibid.* **11**, 1 (1953).
 66. See faunal lists in L. S. B. Leakey, *Olduvai Gorge* (61).
 67. Other riverine situations have provided selected assemblages of pebble tools referred to the Pre-Chelles-Acheul industries. The Kafuan industry, generally regarded as the oldest such manifestation, has been recovered from very high-level gravels of the Kafu and Kagera rivers in western and southeastern Uganda, respectively [C. van Riet Lowe, *Mem. Geol. Survey Uganda* No. 6 (1952); *Proc. Pan-African Congr. Prehist.*, 3rd Congr., Livingstone, 1955 (1957), p. 207]. The industry, in a more evolved facies it seems, is probably also represented in the Katanga, southeastern Belgian Congo [G. Mortelmans, "Mélanges en hommage au Prof. Hamal Nandrin," *Publ. soc. roy. Belge anthropol. préhist.* (1952), p. 150; *Actes congr. panaf. préhist.*, 2^e Congr., Alger, 1952 (1955), p. 295; *Proc. Pan-African Congr. Prehist.*, 3rd Congr., Livingstone, 1955 (1957), p. 214] and in the calcified basal older gravels of the Vaal river in southern Africa [C. van Riet Lowe, *S. African Archaeol. Bull.* **8**, 27 (1953)]. There is little question that some of the selected specimens are indeed artifacts of a very primitive stone industry although this is likely not true of those from the Kafu river; until larger assemblages are found *in situ*, preferably on old land surfaces, in clays or other fine sediments, there will always be a considerable measure of doubt concerning their full authenticity.
 68. J. T. Robinson, *Am. J. Phys. Anthropol.* **12**, 181 (1954).
 69. S. L. Washburn and B. Patterson, *Nature* **167**, 650 (1951).
 70. H. B. S. Cooke, *S. African Archaeol. Bull.* **7**, 59 (1951); F. C. Howell, *Am. J. Phys. Anthropol.* **13**, 635 (1955); R. F. Ewer, *S. African Archaeol. Bull.* **9**, 41 (1956); *Proc. Pan-African Congr. Prehist.*, 3rd Congr., Livingstone, 1955 (1957), p. 135.
 71. K. P. Oakley, *Am. J. Phys. Anthropol.* **12**, 9 (1954); *Proc. Pan-African Congr. Prehist.*, 3rd Congr., Livingstone, 1955 (1957), p. 155.
 72. Sterkfontein, Swartkrans, and Kromdraai are situated in a shallow valley about 6 miles northwest of Krugersdorp, southern Transvaal; Limeworks Cave is located in the Makapansgat valley, several miles northeast of Potgietersrus, central Transvaal; the limeworks at Buxton, near Taung, is just west of the Transvaal border, adjacent to the Kaap escarpment bordering the Harts River, in Bechuanaland (Cape Province).
 73. C. K. Brain, *Proc. Pan-African Congr. Prehist.*, 3rd Congr., Livingstone, 1955 (1957), p. 143; and *Transvaal Museum, Pretoria, Mem. No. 11* (1958), p. 1; also, H. B. S. Cooke, *S. African J. Sci.* **35**, 204 (1938); S. H. Haughton, *Trans. Geol. Soc. S. Africa* **50**, 55 (1947).
 74. F. E. Peabody, *Bull. Geol. Soc. Am.* **65**, 671 (1954); also, R. B. Young, *Trans. Geol. Soc. S. Africa* **28**, 55 (1925).
 75. R. A. Dart, *Transvaal Museum, Pretoria, Mem. No. 10* (1957), p. 1; *Proc. Pan-African Congr. Prehist.*, 3rd Congr., Livingstone, 1955 (1957), p. 161; *Am. Anthropol.* **60**, 923 (1958).
 76. The recent account by W. I. Eitzman [*S. African J. Sci.* **54**, 177 (1958)] of the original constitution of the Limeworks Cave and the mining activities which largely destroyed it suggests that there were probably three separate fossiliferous breccias, separated by travertines in the lower portion of this great cavern. He states that the lower and middle of these were densely packed with bone, whereas the upper horizon was more discontinuous and far less fossiliferous. It is clear that only a very small proportion of the bone accumulations have been salvaged from the mine dumps and that the great bulk of the fossil mammals (and australopithecines) were consumed in the lime kilns.
 77. J. T. Robinson, *Ann. Trans. Museum, Pretoria* **22**, 1 (1952).
 78. R. A. Dart, *Intern. Anthropol. Ling. Rev.* **1**, 201 (1953); *Smithsonian Inst. Ann. Rept.* 1953 (1956), p. 317; *Am. Anthropol.* **60**, 715 (1958).
 79. H. Breuil, *Palaeontol. Sinica, Peking*, **D6**, 1 (1939).
 80. S. L. Washburn, *Am. Anthropol.* **59**, 612 (1957).
 81. A. R. Hughes, *S. African J. Sci.* **51**, 156 (1954); *Am. J. Phys. Anthropol.* **12**, 467 (1954); R. A. Dart, *Am. Anthropol.* **58**, 40 (1956).
 82. J. Stevenson-Hamilton, *Wild Life in South Africa* (Cassell, London, 1954).
 83. R. A. Dart, *Nat. Hist.* **26**, 315 (1926); *Am. J. Phys. Anthropol.* **7**, 1 (1949); see also, K. P. Oakley, in *Appraisal of Anthropology Today*, S. Tax et al., Eds. (Univ. of Chicago Press, Chicago, 1953), pp. 28-30.
 84. C. K. Brain et al., *Nature* **175**, 16 (1955); see K. P. Oakley, *Antiquity* **30**, 4 (1956). This horizon has also yielded a fragment of an australopithecine maxilla [R. A. Dart, *Nature* **176**, 170 (1955)].
 85. J. T. Robinson and R. J. Mason, *Nature* **180**, 521 (1957); see K. P. Oakley, *Antiquity* **31**, 199 (1957). Foreign pebbles were first reported from this region (at Kromdraai) by S. H. Haughton, *S. African Archaeol. Bull.* **2**, 59 (1947).
 86. J. T. Robinson, *Leech* **28**, 94 (1958).
 87. J. D. Clark, personal communication.
 88. F. Weidenreich, *Anat. Anz.* **44**, 497 (1913); A. Keith, *Brit. Med. J.* **1**, 451, 499, 545, 587, 624, 669 (1923); S. L. Washburn, *Cold Spring Harbor Symposia Quant. Biol.* **15**, 67 (1950); L. W. Mednick, *Am. J. Phys. Anthropol.* **13**, 203 (1955).
 89. R. A. Dart, *ibid.* **7**, 301 (1949); *ibid.*, in press.
 90. R. Broom et al., *Transvaal Museum, Pretoria, Mem. No. 4* (1950), p. 1; R. Broom and J. T. Robinson, *Am. J. Phys. Anthropol.* **8**, 489 (1950).
 91. —, *Transvaal Museum, Pretoria, Mem. No. 6* (1952), p. 1.
 92. W. E. Le Gros Clark, *Am. J. Phys. Anthropol.* **13**, 19 (1955); F. C. Howell, in preparation.
 93. W. E. Le Gros Clark, *J. Roy. Anthropol. Inst.* **80**, 37 (1952); J. T. Robinson, *J. Dental Assoc. S. Africa* **7**, 1 (1952); *Transvaal Museum, Pretoria, Mem. No. 9* (1956), p. 1.
 94. —, *Evolution* **8**, 324 (1954); *Z. Morphol. Anthropol.* **46**, 269 (1954); also, see the excellent summary by G. Heberer, *Primates*, I (1956), p. 379.
 95. —, *Am. J. Phys. Anthropol.* **11**, 445 (1953).
 96. See G. H. R. von Koenigswald, *Proc. Koninkl. Ned. Akad. Wetenschap.* **B59**, 318 (1956). This author would distinguish *Paldipithecus rhenanus*, of which a complete right femur is known, from the Pontian of Rheinhesse and the Schwäbian lignites, giving it as a separate genus (rather than as a species, as I have done here).
 97. W. E. Le Gros Clark and D. P. Thomas, *Fossil Mammals of Africa* (British Museum of Natural History, London, 1951) vol. 3, p. 1; H. Zapfe, *Am. J. Phys. Anthropol.*, in press; J. Hürzeler, *Ann. paléontol.* **40**, 1 (1954).
 98. W. E. Le Gros Clark, *Quart. J. Geol. Soc. London* **105**, 225 (1950); — and L. S. B. Leakey, *Fossil Mammals of Africa* (British Museum of Natural History, London, 1951), vol. 1, p. 1.
 99. W. E. Le Gros Clark, *Proc. Zool. Soc. London* **122**, 273 (1952); J. Napier, *Fossil Mammals of Africa* (British Museum of Natural History, London, in press).
 100. H. de Terra, *Science* **124**, 1282 (1956).
 101. See A. Remane, *Abh. Akad. Wiss. Lit. Mainz, Math.-Naturw. Kl.*, No. 2 (1955).
 102. J. Hürzeler, *Schweiz. Palaeontol. Abhandl.* **66**, 1 (1949); *Eclogae Geol. Helv.* **44**, 404 (1951); *Verhandl. naturforsch. Ges. Basel* **65**, 88 (1954); *ibid.* **69**, 1 (1958).
 103. See W. L. Strauss, Jr., *Quart. Rev. Biol.* **24**, 200 (1949).
 104. In Mediterranean Africa the Pontian is little known in southern and northern Tunisia and practically not at all in Morocco. The richest locality, except for primates, is Qued el Hamman, Oran (Algeria). Findings from the later Pliocene are extremely rare; this period is known largely from the Constantine (Algeria) sites of Ain el Bey and Ain el Hadj Baba. A fairly substantial faunal assemblage, probably middle Pliocene, from the Gart el Moluk, Wadi Natrun, in Egypt, is known. The situation is even more discouraging south of the Sahara where a Pliocene fauna is known only from diamond diggings in Little Namaqualand, southwest Africa.
 105. L. Cahen and J. Lepersonne, *Mem. soc. géol., paléontol., hydrol. Belg. ser. 8*, No. 4 (1952), p. 1; J. de Heinzelin, "Soils, paléosols et désertifications anciens," *Publ. Inst. Natl. Etude Agron. Congo Belge, Bruxelles* (1952).

High-Altitude Observation Techniques

A variety of tools besides satellites are available for the study of upper atmosphere physics.

Raymond C. Staley

It may be surprising to many not acquainted with upper atmosphere research to learn that the greater part of all we know today about the structure of the upper atmosphere has been discovered without instrumented satellites. Two important bits of information provided by satellites about the upper atmosphere have so far been made public: (i) The atmosphere from 500 to 1000 km above the earth is between 16 and 40 times as dense as was thought in 1955. (ii) The soft radiation, previously detected above 50 km by balloon- and rocket-borne Geiger counters and now known to be contained in two belts at about 3000 and 14,000 km, is at least 100 times as great as was formerly believed (1). This survey has been prepared to explain briefly how the chemical composition, ionization, density, temperature, and motion of the atmosphere above 10 km can be studied by methods some of which use balloon and rocket instrumentation but most of which actually use ground-based observations of atmospheric phenomena.

Some Methods of Study

A classical method used to give good estimates of upper air temperatures and winds involves the anomalous propagation of sound. This method presupposes a knowledge of the molecular weight and the ratio of specific heats for the atmosphere at all levels traversed by the sound wave. The maximum height at which the wind and temperature can thus be measured is usually taken as about 50 to 60 km for ground-based explosions. With grenades exploded from rockets, the upper limit may be extended to 80 km (2). Sound waves with very low frequencies

(0.1 cy/sec) can be bent back in the ionosphere at altitudes of about 170 km (3). Such sound signals can give a value for the expression

$$c_p RT/c_v m$$

where c_p/c_v is the ratio of specific heats and R/m is the gas constant for the medium. Unless these ratios are known, the temperature T ($^{\circ}\text{K}$) cannot be found. According to Hulbert (4) the composition of the atmosphere is approximately constant to about 150 km, and, hence, the use of sound waves is valid for the study of temperature and wind within this region.

Winds, temperature, atmospheric composition, solar radiation, and electrical conductivity have been measured with instruments sent as high as 42 km with large balloons (5). Similarly, rockets have been instrumented for studies at still higher altitudes. Mass spectrographs and ion spectrographs carried aloft have provided information about the presence of argon and of atomic and molecular forms of both oxygen and nitrogen up to 220 km. Photon counters sent aloft have measured solar ultraviolet and x-ray energy available for ionization and photodissociation (1, 4). In addition, other instrumentation has verified the indirectly determined vertical variation of temperature and has given much information about cosmic and soft radiation (1).

Before the atmosphere beyond the range of balloons is considered, some mention might be made about the study of the vertical distribution of atmospheric ozone with ground-based instruments. Götz, Dobson, and others (6) found means of determining the total amount and the vertical distribution of ozone. The total amount is determined

by measuring the intensity of solar radiation at two different wavelengths, one at which the ozone absorption is strong, a second at which it is weak. The measurements are made at two different zenith angles. With proper measurements and laboratory-determined constants, the total ozone can be found. The mean height of the ozone layer may be estimated by studying the ozone absorption in light scattered from the zenith with the sun at various low elevations (7). Epstein (8) has presented a ground-based method for finding the vertical ozone distribution from knowledge of the total ozone content together with knowledge of the infrared absorption of solar radiation by ozone and the rate at which ozone radiates in the infrared. Paetzold (9) reports that a study of the sunlight reflected from the moon's surface right at the edge of the earth's shadow during a lunar eclipse will allow a determination of ozone distribution in the earth's atmosphere up to 45 km for various latitudes. The argument is that with a tangential path of solar rays there is a large decrease in the yellow-orange (Chappuis) spectral region, causing a blue-green coloring of the earth's shadow near the shadow limit. This zone represents an optical copy of the atmospheric ozone layer.

An interesting ground-based technique for finding the density distribution in the atmosphere between 10 and 70 km is the nighttime use of a modulated searchlight beam directed upward. The back-scattered light is observed from the ground at a point about 20 km from the searchlight. If Rayleigh scattering is assumed, the intensity of light will be a function of air density (10).

The observation of meteorites both by photographic means and by radio echoes provides considerable information about the atmosphere from 60 to 120 km. Air density, winds, and wind shear have been deduced from photographs of meteorites (11). Booker (12) states that information about atmospheric turbulence can be deduced from photographic observation of meteor trails. Atmospheric pressures and scale heights can be found from radio studies of meteor trails (13). Robertson (14) reports that a 27-megacycle-per-second

The author is a research meteorologist at the electrical engineering research laboratory and assistant professor of meteorology in the department of aero-space engineering, University of Texas, Austin. He is spending the current academic year at the department of meteorology and climatology, University of Washington, Seattle, on a National Science Foundation faculty science fellowship.

Doppler-radio system has been used to estimate winds from meteor trail drift at 80 to 105 km; these winds appear to have prominent 12- and 24-hour tidal harmonics. Greenhow (15) has calculated the atmospheric diffusion coefficient by use of 8.27- to 8.35-meter radio measurements of echoes from meteor trails between 80 and 100 km.

Winds in the region between 30 and 90 km have been measured both by the observation of sound waves from grenades released from rockets (2, 16) at various altitudes and by the observation of radar return from vaporized, metal-coated, thin plastic strips (chaff) released from rockets (17). Metallic sodium vapor ejected from Aerobee rockets during twilight hours creates a sodium emission cloud at 5890 angstroms, and observation of these clouds provides wind information at 85 to 200 km (18, 19). Figure 1 shows wind-distorted sodium trails extending from an altitude of 100 km to about 200 km. Methods for computing wind velocities from such photographs have been described by Manring (19).

The natural phenomena known as noctilucent clouds, which occur at heights of about 80 km and apparently consist of volcanic or meteoric dust, offer some opportunities for making qualitative estimates of air turbulence and vertical stability. Ludlam (20) states that theodolite tracking of a noctilucent cloud element to get its velocity is difficult because these clouds usually lack elements which are well defined and which retain their identity. Rapid changes in the clouds and the presence of billows suggest a steep lapse rate and considerable stirring. Noctilucent clouds probably do not exist at latitudes less than 55 degrees, and so this source of information is limited geographically.

Ionospheric Studies

Studies of the ionosphere have provided information about the electron density and the temperature in the ionized layers; attempts have also been made to measure the size as well as the drift of the ionospheric irregularities. Radio noise should contain a component due to thermal radiation from the ionosphere, and this radiation, which is very weak, has been identified and measured (21); temperatures measured in this way range from 240° to 290°K. The rate of ion production at a given height is extremely sensitive to tem-

perature changes, and so diurnal and seasonal height changes in ion-forming layers are found to correspond to diurnal and seasonal temperature changes (22). The daily and annual variations in F layer ionization (at about 180 km) suggest that the gradient of the temperature inversion in that layer is less in summer than in winter, and also that the warm layer is thicker in summer than in winter (23). Studies of the E and F₂ layers, with ionization ascribed to atomic oxygen, give temperatures of 190° to 220°K at about 110 km and 500° to 1200°K at about 250 km (24). The wide range is partly seasonal and partly related to solar activity, with the

highest temperatures occurring in a summer of great solar activity.

Observation of radio echoes reflected from the moon shows slow variations in the mean daytime amplitude of the echoes. The plane of polarization of the radio waves is rotated due to the presence of the earth's magnetic field, and the extent of rotation depends upon the total electron content along the line of sight to the moon. J. V. Evans (25) explains that a technique has been developed for the deduction of the amount of rotation and thus for the determination of the total electron content.

Scintillations of radio stars have been shown to be caused by phase changes




Fig. 1. Two sets of simultaneous photographs of a morning-twilight sodium trail, New Mexico, 26 November 1957. The upper pair, showing the trail over an altitude range from about 110 to 200 km, were taken when the rocket was near the zenith; the lower pair were taken about 5 minutes later. The left-hand pictures were made at Socorro, 211 km north-northeast of Deming, where the right-hand pictures were taken. [Courtesy Air Force Cambridge Research Center]

in the incoming radiation as it traverses irregularities in the electron density of the F region. The motion of ionization in the F region is probably related to turbulence and eddies in the dynamo region (135 to 150 km) through interaction between the earth's magnetic field and the electric field (26).

Vestine (27) has shown that, since geomagnetic disturbances can be represented by atmospheric electric-current systems and, in turn, the current systems can be replaced by equivalent atmospheric wind systems, it may be possible to deduce air motion in the E and F layers (100 to 300 km) from geomagnetic data.

Benyon and Goodwin (28) note that at about 100 km there is a high correlation between the magnitude of the drift velocity of ionospheric irregularities and the horizontal temperature gradient. This suggests that the study of the vertical structure of ionospheric drift may provide important information about the temperature of this part of the atmosphere.

Density beyond 300 Kilometers

Recent spectrographic and radio studies of the solar corona seem to support the idea that the atmosphere of the sun extends to and beyond the earth. According to Chapman (29) the earth and its atmosphere are immersed in this solar gas and must act as a sink of heat. Nicolet (30) suggests a figure for the heat conducted by atomic hydrogen from the corona to atomic oxygen in the earth's atmosphere of about 36 ergs per square centimeter per minute (as compared with the "solar constant" of about 8.4×10^7 erg/cm² min). Chapman's argument finds support in the comparatively large densities discovered from tracking artificial satellites and also from studying radio whistlers. Whistlers apparently are caused by thunderstorm electrical discharge (sferics) with an energy spectrum peaking in the 10- to 20-kilocycle-per-second range. The sferics travel through the ionosphere and on to altitudes of 500 to 5000 km or more, following the lines of flux of the earth's magnetic field out and then back to the opposite hemisphere. For this to occur, the average electron density of the atmosphere must be considerably greater than any previous estimates, about 400 electrons per cubic centimeter (31). Studies of the extension of the solar corona can be made by observing how radio waves

from the radio star Taurus vary in intensity as they travel the outer corona (32). Since anomalies in the solar gas cloud appear to be related to changes in the earth's magnetic field, it may be possible to learn more about these anomalies by means of induced currents in transoceanic cables (1).

The flow of atomic hydrogen into the outer atmosphere means that the outer atmosphere must be more dense than has been thought. It is interesting to note that one piece of indirect evidence led for a time to contrary conclusions. From studies of the absorption of solar x-rays and the dissociation of oxygen up to about 165 km, some observers, by extrapolating the derived density distribution, calculated an upper atmosphere of considerably smaller density (33). This view was bolstered for a time by the idea (reminiscent of the early arguments about the stratosphere) that the high atmosphere should be isothermal and in radiational equilibrium. Chapman and Nicolet have countered such arguments, and the weight of evidence is for the more dense upper atmosphere above 300 km, with the temperature, as represented by the kinetic energy of the molecules, increasing with height. Here we have a case of different indirect evidence giving far different ideas about the density of the upper atmosphere. On the one hand the arguments for thermal equilibrium called for a low-density, isothermal upper atmosphere, while the observations of whistlers called for a contrary conclusion. Finally, a third indirect method, by visual, photographic, and radio tracking of the artificial satellites, provided evidence in support of the denser atmosphere. Sterne (34) gives a simplified formula for inferring air density at perigee from visual observations of artificial satellites. Calculations based on observations of sputnik I and reported by Harris (35) give an atmosphere at 400 km which is 40 times as dense as the ARDC (Air Research and Development Command) model atmosphere at that level. Later data, reviewed by Kallman (36), are in substantial agreement with Harris' conclusions.

A device constructed but not yet used for determining high-altitude density is the satellite balloon, a 12-foot, spherical plastic, aluminum-covered balloon carried aloft in collapsed state by an Explorer satellite. When the satellite is in orbit, the balloon is released and inflated by a bottle of gas. It then will orbit like the satellite, but air drag will act to slow the balloon

more rapidly than it slows the satellite. A study of the positions of the satellite and balloon should give more information about the density of the outer atmosphere. The balloon would be large enough to be visible by eye from the earth's surface during twilight hours (37).

Studies of the emission spectrum of the aurora and night airglow provide information about constituents of the outer atmosphere. The aurorae are most frequent at heights of 100 to 120 km, but they are observed from 63 to 400 km. Sunlit aurorae may occur as high as 1100 km (38, p. 422). Atomic oxygen, nitrogen, and hydrogen, the oxygen and nitrogen ions, O₂⁺ and N₂⁺, and molecular oxygen and nitrogen have all been observed in aurorae. Seaton (39) has estimated the N₂/O ratio for heights between 160 and 400 km. He also gives an N₂⁺ density above 740 km as greater than 120 ions per cubic centimeter. The night airglow emission is known to originate at heights from 70 km (OH) to 1000 km (oxygen) (38, p. 533). Its spectrum, which shows atomic oxygen and sodium, the N₂⁺ ion, molecular oxygen and nitrogen, and the OH radical, runs from the violet well into the infrared (10 μ). At one time it was thought that the infrared emission was caused by the recombination of atomic nitrogen, but this emission is now attributed to vibration-rotation bands of hydroxyl (40). Phillips (41) has observed the width of the oxygen emission line 5577 Å in the airglow and the line 6300 Å in the twilight flash. These line widths indicate temperatures between 155° and 231°K at 100 km for the former and a temperature of about 750°K for the latter at about 220 km. Studies of the translation and rotation of airglow cells (5577 Å emission) have convinced personnel at the University of Colorado High Altitude Observatory that these motions represent true winds at 100 km (42).

Conclusions

From this brief survey of some methods available for upper atmosphere research, it appears that there are many tools of widely varied complexity which can be used for such work. Most of these need further research and development. It would seem wise to use the most promising of the indirect measurement techniques near rocket-launching areas and to make the measurements synoptic with rocket studies. It is also

important that measurements of the outer ionosphere be matched with satellite data. It is not unlikely that, once certain relations are verified, ground-based observations will be capable of providing hour-by-hour and day-by-day information about the upper atmosphere which would be much too costly to obtain routinely with rockets. Furthermore, some of the indirect methods will permit routine measurements from regions too close to the earth for observations with long-lived satellites.

References

1. H. Odishaw, *Science* **128**, 1599 (1958).
2. C. J. Bracefield, *J. Geophys. Research* **59**, 233 (1954).
3. J. Veldkamp, *J. Atmospheric and Terrest. Phys.* **1**, 147 (1951).
4. E. O. Hulbert, *Meteorol. Monographs* **3**, 160 (1957).
5. R. J. Murgatroyd, *Quart. J. Roy Meteorol. Soc.* **83**, 424 (1957).
6. R. M. Goody, *Physics of the Stratosphere* (Cambridge Univ. Press, Cambridge, England, 1954), p. 81.
7. F. W. P. Götz, "Ozone in the atmosphere," in T. F. Malone, *Compendium of Meteorology* (Am. Meteorol. Soc., Boston, Mass., 1951), p. 275.
8. E. S. Epstein, C. Ostergard, A. Adel, *J. Meteorol.* **13**, 319 (1956).
9. H. K. Paetzold, *J. Geophys. Research* **59**, 365 (1954).
10. L. Elterman, "Seasonal trend of temperature, density and pressure in atmosphere obtained with searchlight-probing technique," *U.S. Air Force Cambridge Research Center Geophys. Research Paper No. 29* (1954); S. S. Frieland, J. Katzenstein, M. R. Zatzick, *J. Geophys. Research* **61**, 415 (1956).
11. F. L. Whipple, *Advances in Geophys.* **1**, 119 (1952).
12. H. G. Booker, *J. Atmospheric and Terrest. Phys. Suppl.*, **52** (1957).
13. S. Evans, "Atmospheric pressures and scale height from radio echo observations of meteors," in T. R. Kaiser, *Meteors* (Pergamon Press, London, 1955), p. 86.
14. D. S. Robertson, D. T. Liddy, W. G. Elford, *J. Atmospheric and Terrest. Phys.* **4**, 255 (1953).
15. J. S. Greenhow and E. J. Neufeld, *ibid.* **6**, 133 (1955).
16. W. G. Stroud, W. Nordberg, J. R. Walsh, *J. Geophys. Research* **61**, 45 (1956).
17. L. B. Smith, *Bull. Am. Meteorol. Soc.* **39**, 436 (1958) (abstr.).
18. H. D. Edwards, "Emission from a sodium cloud artificially produced by means of a rocket," in E. B. Armstrong and A. Dalgarno, *The Airglow and the Aurorae* (Pergamon, London, 1956), p. 122.
19. E. Manring, J. F. Bedinger, H. B. Pettit, *J. Geophys. Research* **64**, 587 (1959).
20. F. H. Ludlam, *Tellus* **9**, 341 (1957).
21. J. L. Pawsey, L. L. McCready, F. F. Gardner, *J. Atmospheric and Terrest. Phys.* **1**, 261 (1951).
22. D. Lepechinsky, *ibid.* **1**, 278 (1951).
23. O. Burkhard, *ibid.* **8**, 83 (1956).
24. S. S. Baral and A. P. Mitra, *ibid.* **1**, 95 (1950).
25. J. V. Evans, *ibid.* **11**, 259 (1957).
26. M. Dagg, *ibid.* **11**, 133 (1957).
27. E. H. Vestine, *J. Geophys. Research* **59**, 93 (1954).
28. W. J. G. Benyon and G. L. Goodwin, *J. Atmospheric and Terrest. Phys.* **13**, 180 (1958).
29. S. Chapman, "Achievements and prospects in auroral and airglow research," in E. B. Armstrong and A. Dalgarno, *The Airglow and Aurorae* (Pergamon, London, 1956), p. 1; "Notes on the solar corona and the terrestrial ionosphere," *Smithsonian Contribs. to Astrophys.* **2**, No. 1 (1957).
30. M. Nicolet, *Science* **127**, 1317 (1958).
31. "IGY Bulletin," *Trans. Am. Geophys. Union* **38**, 1011 (1957).
32. A. Hewish, *Proc. Roy. Soc. London A228*, 238 (1955).
33. H. Wexler, *Trans. Am. Geophys. Union* **38**, 954 (1957).
34. T. E. Sterne, *Science* **127**, 1245 (1958); T. E. Sterne and G. F. Schilling, "Some preliminary values of upper atmosphere density from observation of U.S.S.R. satellites," *Smithsonian Contribs. to Astrophys.* **2**, No. 10, 207 (1958).
35. I. Harris and R. Jastrow, *Science* **127**, 471 (1958).
36. H. K. Kallman, *J. Geophys. Research* **64**, 615 (1959).
37. News stories, *New York Times*, 29 Mar. and 13 Apr. 1958.
38. S. K. Mitra, *The Upper Atmosphere* (Asiatic Society, Calcutta, India, 1952).
39. M. J. Seaton, "Excitation processes in the aurora and airglow," in E. B. Armstrong and A. Dalgarno, *The Airglow and the Aurorae* (Pergamon, London, 1956), p. 225.
40. V. I. Drassovsky, "On the detection of the infrared night airglow," *ibid.*, p. 86.
41. J. G. Phillips, "The determination of the widths of the airglow and the twilight flash," *ibid.*, p. 67.
42. F. E. Roach, E. Tandberg-Hanssen, L. P. Megill, *J. Atmospheric and Terrest. Phys.* **13**, 122 (1958).

Science in the News

Khrushchev Sees Aspects of American Science and Talks About Soviet Achievements

Soviet Premier Nikita S. Khrushchev's trip to the United States closed further the narrowing gap between science and politics. In his statement on arrival, the Premier made the first of many scientific references when he spoke of the Soviet moon rocket and of his country's newly launched atomic icebreaker. Of the moon rocket, he said:

"Shortly before our meeting, Mr. President, the Soviet scientists, engineers, technicians and workers filled our hearts with joy by launching the

rocket to the moon. . . . A container . . . with a pennant bearing the national emblem of the Soviet Union is now on the moon. . . . We entertain no doubt that the splendid scientists, engineers and workers of the United States of America who are engaged in the field of conquering the cosmos will also carry their pennant over to the moon. The Soviet pennant, as an old resident of the moon, will welcome your pennant, and they will live there together in peace and friendship as we both should live together on the earth. . . ."

The atomic icebreaker that was mentioned is the 16,000-ton *Lenin*, which started its maiden voyage on 15 September, the day that Mr. Khrushchev

arrived in the United States for his 12-day stay. The ship, which can carry enough fuel to cruise for several years, is designed to keep open the 11,000-mile arctic sea route between Murmansk and Vladivostok—a route at present open about 10 weeks each year.

Disarmament Plan Offered

Disarmament, including a nuclear-test agreement and cooperation in the peaceful use of atomic energy and outer space, was a persistent theme in virtually all of the Soviet Premier's speeches. He set the stage for his total disarmament proposal to the United Nations when he addressed the National Press Club in Washington on the second day of his visit. He said:

"The best, the most reliable way to make war impossible would be to place all states, without exception, in conditions where they would have no means of conducting war. . . ."

"The Soviet Union and the United States are faced with this alternative: Either the latest achievements of scientific and technical thought—the discovery of the secret of the atom, the development of rockets and the penetration of outer space—will be placed

in the service of a peaceful future and prosperity of mankind, or they will be used for the purpose of destruction and annihilation and, as a result, the earth will be covered with ashes and graves. The Soviet people have long made their choice for peace."

Next day he repeated these views in a general discussion of peaceful coexistence before the members of the Senate Foreign Relations Committee and guest senators.

During an hour-long question period, Senate Majority Leader Lyndon Johnson (D-Tex.) asked whether the Soviet Union is willing to exchange secrets on space flight. Mr. Khrushchev replied that he does favor such exchange. When Johnson pressed him then as to why the Soviet Union had not participated in the U.N. *ad hoc* committee on this subject, he said that the United States had attempted to put the Soviet Union into the position of a "poor relation," that the Soviet Union had not been treated with the proper respect in this area, and that it was a subject on which the two powerful countries active in the field could agree.


On the following day, 18 September, the Soviet Premier gave the United Nations his 4-year plan for general and complete disarmament of all states. Some of the scientific aspects of the plan are as follows:

"All atomic and hydrogen bombs . . . will be destroyed and their further production terminated. The energy of fissionable materials will be used exclusively for peaceful economic and scientific purposes.

"Military rockets of all ranges will be liquidated and rocket facilities will remain only as means of transportation and harnessing outer space. . . . The stockpiles of chemical and bacteriological means of warfare accumulated by some states . . . will be eliminated once and for all. . . ."

Later in the address Khrushchev considered control, proposing the establishment of an international body to set up a system of control over all disarmament measures. After first stating that the Soviet Government would not resume nuclear tests if the Western powers do not resume theirs, he said:

" . . . The elaboration of a program of general and complete disarmament should not hold up the solution of such an acute and fully mature question as that of the discontinuance of nuclear weapons tests for all time. There now



Premier Khrushchev visits the Agricultural Research Center in Beltsville, Md., where he hears John W. Mitchell explain how chemicals can regulate plant growth and examines a cow especially bred for high milk production.

exist all prerequisites for such a solution. We hope that the appropriate agreement on the discontinuance of tests will be concluded and put into effect without delay."

Initial reaction in the United States to the disarmament proposals was restrained. Most official observers were reluctant to comment, except to suggest that at important points the plan was vague. Interest was expressed, however, in Mr. Khrushchev's test-ban remarks. Some observers, such as Senator Hubert H. Humphrey, chairman of the Senate Subcommittee on Disarmament, felt that these comments offered an unparalleled opportunity to initiate a disarmament control system. Others, such as Senator Clinton D. Anderson, chairman of the Joint Committee on Atomic Energy, have indicated that it would be wiser to attempt only a limited agreement, one which would permit continued testing underground without inspection.

The interrelationship of science and public affairs was demonstrated not only by words but also by Soviet Premier Khrushchev's itinerary, which listed an impressive number of research laboratories and agricultural and technical installations.

Beltsville Visited

Because increasing Soviet food production is a major aspect of the U.S.S.R.'s national policy, and because Mr. Khrushchev was in charge of Russian agriculture before he became Premier, it is not surprising that one of his first actions was to inspect the Department of Agriculture's Agricultural Research Center at Beltsville, Md. He paid a long visit there on 16 September.

In a slightly barbed welcoming speech, Secretary of Agriculture Ezra Taft Benson praised the progress of American agriculture "under our capitalistic free-enterprise system." He noted the cooperation in farm research between federal and state specialists and those in private industry, and called the family farm the "backbone of American agriculture."

The Secretary then turned the program over to some of the department's scientists in the Bureau of Plant Industry. Harry A. Borthwick recently helped discover a form of plant pigment that controls growth. He described how plant growth can be promoted or inhibited through the pigment by exposure of seed and plants to light. Next, Warren C. Shaw discussed a new weed kill-

er that does not harm cereal crops and legumes. John W. Mitchell, shown in one of the accompanying pictures, displayed some results of the application of growth-regulating chemicals to plants, fruits, and trees.

After the lectures, Mr. Khrushchev told Secretary Benson that he thought the Soviet Union could benefit from American agricultural research. He also suggested that there should be an increase in the exchange of technicians.

The last part of the Beltsville visit was spent inspecting the experimental livestock, about which the Premier exhibited considerable knowledge. He is shown examining a cow which, during a 305-day lactation, produced 18,000 pounds of milk and 724 pounds of butterfat. Her forebear, eight generations back, produced only 15,800 pounds and 540 pounds, respectively.

Mr. Khrushchev saw the practical application of scientific farming on 23 September when he went to Iowa to inspect Roswell Garst's hybrid-corn farms and experimental growing plots. The Premier paid particular attention to chemical fertilizers, synthetic feeds, and new machinery. Garst's hybrid seed was exported to the U.S.S.R. in great quantity and opened the way for Soviet introduction of modern corn-farming techniques.

The Premier concluded the agricultural aspect of his tour with a visit to Iowa State College, Ames, a major agricultural research center.

Research Secrecy Discussed

While he was in California, Mr. Khrushchev toured the International Business Machine's computer plant, where he gave some of the workers souvenir medals commemorating the Soviet moon rocket. Later, when he discussed his IBM visit at a joint dinner of the San Francisco Commonwealth Club and the World Affairs Council, he turned to the topic of secrecy in research, saying:

"The plant we saw was making computers . . . I don't know who makes the better machines. . . . For the time being we're keeping them a secret. . . . Both sides are actually keeping them a secret for the time being. . . . But I am sure that the time will very soon come when there will be no such secrets and people will be able to see all such machines and benefit by them."

Two Major Visits Canceled

During the California stay, Mr. Khrushchev was also scheduled to see the

Stanford Research Institute at Menlo Park, but time would not permit and he sent three delegates instead, including Vasily S. Yemelyanov, chief of the Main Administration for the Use of Atomic Energy. The projects shown to the Soviet officials included such subjects as research aimed at finding chemicals to cure cancer, efforts to improve the use of radar to give early hurricane warnings, a machine that uses principles of television to give a high-speed facsimile reproduction process, and research on opportunities for the private investment of capital in foreign countries.

Mr. Yemelyanov is reported to have found it difficult to understand that the institute was organized to sell research services in any field wanted by governmental or private customers. He commented that in the Soviet Union, research is organized for a specific purpose, such as the creation of better airplanes or the conquest of a disease such as cancer.

A second major scientific visit that the Premier was forced to cancel was that to the National Institutes of Health in Bethesda, Md., where an elaborate 1½-hour program had been prepared. A. M. Markov, chief of the Soviet Medical Service, and Dr. V. N. Butrov, of the Soviet Foreign Ministry, went in his place. Before leaving, Markov commented that American and Soviet medical people should join in eliminating disease in all countries:

"Our two countries are more or less equal economically. And if our doctors could get together, we could do very much for the peoples of all nations."

Next day, on 26 September, H. van Zile Hyde, Assistant Surgeon General for International Affairs, announced that the United States and the Soviet Union have agreed to engage in joint health-research projects in cancer, heart disease, and poliomyelitis. This group of projects is the first major cooperative effort in peaceful scientific pursuits to be announced as a result of the Premier's visit to this country.

Mr. Khrushchev's tour of the United States was set in historical perspective by the French paper, *Paris-Presse*, which quoted a comment Lenin made in 1920 to H. G. Wells. Lenin had just read one of Wells' science-fiction books, and is reported to have said:

"I, too, understand that all human conceptions are on the scale of our planet. They are based on the assumption that technical potentials, when

developed to the full, will not go beyond 'terrestrial limits.' If we arrive at establishing interplanetary communications we must revise all our philosophical, social, and moral conceptions. In that case the technical potentials, having become unlimited, will impose upon us the end of violence as a means and method of progress."

House Committee Questions Adequacy of Manpower Roster

In a report last month, the House Science and Astronautics Committee, headed by Overton Brooks (D.-La.), expressed reservations about the completeness and currency of the National Science Foundation's Register of Scientific and Technical Personnel. In criticizing the register, the report cited NSF studies that estimated that there are approximately 300,000 persons in the United States who would qualify under a generally acceptable definition of the term *scientist*. The committee juxtaposed this figure to the 170,000 the register now lists. Questions were also raised about the value of dated information on individuals whose profession is characterized by frequent changes of place of employment and field of investigation. Unable to decide whether the register was receiving sufficient support and planning, the committee asked the Science Foundation to report to it next January.

Answers to the committee's questions are currently being worked up for the NSF report in January. The foundation's position is that "very considerable progress" has been made in the register program. In testimony given during the last sessions of the hearings, NSF spokesmen cited the doubling of funds allotted for the program by the foundation for fiscal years 1959 and 1960. New and larger quarters have been acquired for the Records Center in Raleigh, and additional tabulating and microfilming equipment has been installed. Studies are under way on the problem of speeding up the processing of data as they arrive.

A simpler questionnaire is being prepared and plans are being developed to make the register better known and to make its data more readily available to participating scientists through professional societies.

Although the House committee and the Science Foundation have disagreed

on some matters of interpretation concerning the register, the basic attitude is one of cooperation. The committee is aware of the difficulties that the register faces, some of which stem from the way Congress itself operates. For example, the House Committee on Science and Astronautics may want the register expanded, but it does not follow that the House Committee on Appropriations will give the Science Foundation the necessary funds. As Chairman Brooks himself noted, at the very time the spring hearings on NSF's division of scientific manpower and education were being conducted, the House Committee on Appropriations recommended a cut—later restored—in funds for the unit.

Radio Frequencies for Research Studied at Geneva

The allocation of radio frequencies for research in space is one of the major issues now before a 3-month-long conference of the International Telecommunications Union at Geneva. Radio astronomy and space communications, two fields that have expanded greatly since the last ITU conference in 1947, are the particular subjects of study of the 708 delegates at the conference, which began 17 August and will continue until 17 December. The ITU, a U.N. organization through which countries regulate international telecommunications, is using the long session to overhaul the regulations drawn up in 1947. Rapid technological change in recent years and the opening of new fields of communication led to the current sessions.

On the opening date of the conference, the United States delegation presented its position on the problem of frequencies for radio astronomy. After citing the increasing value of the new science's findings, the U.S. group proposed that a world-wide allocation to radio astronomy be made of the frequency band 1400 to 1427 megacycles per second (Mcy/sec). This band, also called the interstellar hydrogen line band, is the major one used in radio astronomy. Another proposal, put forth by the Netherlands delegation, specified a range from 1399 to 1427 Mcy/sec. Virtual agreement among the conferees is reported on the protection of a band at least as wide as that in the U.S. proposal. The U.S. paper also cited measures taken to afford national pro-

tection to radio astronomy observatories in this country.

In another action, the U.S. delegation proposed that six frequencies be set aside for space communications. These communications would be between the earth and satellites, and between satellite and satellite. The frequencies, which would be used for tracking, guidance, and telemetering of data, are: 1700 to 1725, 1825 to 1850, 2275 to 2300, 8300 to 8400, 15,150 to 15,250, and 31,500 to 31,800 Mcy/sec.

Some Opposition

The assignment of radio frequencies for research is not an easy matter, according to reports on the conference. Many conflicts come up that require extended negotiation. In the assignment of frequencies for space communication, for example, some opposition was registered by the delegation from the Soviet Union on the grounds that such allocations at this time would be premature. Pressure from the other delegations moved the Russian group to ask for more time to study the proposals. A number of organizations, particularly military and commercial users, object to the assignment of frequencies for research because the bands might be needed in the future for nonresearch uses. Conference proceedings are aided, however, by a considerable body of accepted practice that has accumulated during the years the ITU has been in existence. Because of this common body of accepted practice, the Union, which was formed by the merger of two of the oldest inter-governmental organizations in the world, generally manages to resolve the conflicts that arise.

Health Research Grant Practices Held Satisfactory

Large increases of funds for the National Institutes of Health will not result in a lowering of the quality of the research projects supported by NIH, spokesmen at the Bethesda, Md., facilities report. The review process for grant requests, and the growing volume of requests for funds for medical research, officials say, assure that the new funds will be used wisely.

Last August, President Eisenhower expressed some concern on this point when he signed the appropriations bill for the Department of Health, Education, and Welfare under which NIH

operates. Although his dissatisfaction was not sufficient to make him veto the bill, which increased NIH's funds from \$294 to \$400 million, he did indicate that he had reservations about the wisdom of a 36 percent increase in funds for a single year. His first concern was that the large increase should "lower the quality of the projects supported by increasing the flow of grant applications more rapidly than the procedures for their careful appraisal can be effectively adapted." He directed the Secretary of Health, Education, and Welfare and the Surgeon General to take appropriate steps to assure themselves that any new research projects be of "high priority and great promise." The President's reservations and his directives to the officials of the department turned the spotlight squarely on the procedures of evaluation that now exist in NIH. Are they adequate now? Will they be changed or expanded?

The NIH grant evaluation process, with its commentary on individual grant requests by study sections, rating of them by advisory councils, and approval by the Surgeon General of as many of the highest-rated projects as the money allows, has been in operation since 1946 when it was devised by C. J. Van Slyke, now Deputy Director of the Institutes. In the opinion of NIH spokesmen, the system has informally been doing the job that the President formally directed the officials of the Department of Health, Education, and Welfare to do in his statement when he signed the appropriations bill. The effect of the growing appropriations for NIH, spokesmen say, is that the Surgeon General's office will be able to go lower in the stack of grant requests. But, because of the very large number of the grant applications and the stringency of the review process, NIH officials feel, standards will not suffer and the new money will go to research projects that will add significantly to the public health.

Congress Leaves Conservation Issues for Next Year

Members of the 86th Congress brought their first session to an adjournment on 15 September and left most of the important conservation legislative issues for the second session that will begin in January. The National Wildlife Federation reports that, although the session did not result in any outstanding conservation accomplishments, several measures were brought along to points

where final action may be expected early next year.

Among the major conservation bills under consideration, that strengthening the Federal Water Pollution Control Act probably made the most progress. This measure to protect streams from raw sewage pollution (H.R. 3610) was sponsored by Representative John A. Blatnik (Minn.). It passed the House, 255 to 143, and won approval in the Senate by a 61-to-27 margin. The House version of the bill, however, would increase the program of grants for the construction of municipal sewage treatment works from \$50 million to \$100 million annually for 10 years, while the Senate voted only \$80 million. Differences in the measures were being considered in a Senate-House conference committee at adjournment time, and the compromise version probably will be among the first sent to the President next year.

Soil Bank and Pesticide Awards

After lengthy discussion, Congress granted \$375 million—the same amount as last year—for the Conservation Reserve Soil Bank program, a program for reducing surplus agricultural production. Under the Soil Bank, farmers lease land to the Federal Government, which uses it for wildlife preserves or other purposes. Last year 23 million acres were retired under this plan at a cost of \$375 million. If this land had been planted, it would have cost the government \$600 million just to store the surplus produced. An even greater controversy over the Soil Bank can be expected in the next session, for the act authorizing the program expires at the end of the 1960 calendar year and must be extended if the plan is to continue. Measures to extend the authorization have been introduced but have not yet received serious consideration.

Congress also approved S. 1575 authorizing the expansion of the Fish and Wildlife Service research program into the effects of pesticides on fish and wildlife. The measure, which contains a maximum authorization of \$2,565,000 per year, was signed by the President on 16 September. Efforts by conservationists to knock out the \$2.4 million Department of Agriculture appropriation for fire ant "eradication" failed, however. This program has a serious detrimental effect on fish and wildlife, and other organisms.

Other Measures Held Over

One of the few major conservation measures not involved in budgetary

considerations—the Wilderness Bill to preserve public lands (S. 1123)—was included in the legislation held over until the second session. Although it has not yet been voted on by committees in either the Senate or the House, much groundwork was laid toward working objections out of the bill and in holding extensive field hearings. Action by the Senate Committee on Interior and Insular affairs is expected early next year.

Among other measures to be held over into the second session are: S. 812, to establish a Youth Conservation Corps, which has passed the Senate; S. 1262, to authorize large impoundment fisheries research, which has passed the Senate; H.R. 2565, the Sikes Bill to authorize fish and game programs on federal military reservations, which has been reported by the House Merchant Marine and Fisheries Committee; H.R. 7045, to establish the huge Arctic Wildlife Range in Alaska, also reported by the House Merchant Marine and Fisheries Committee; and S. 2086, to establish a wildlife disease laboratory, which has been reported from the Senate Interstate and Foreign Commerce Committee.

National Medical Library Interlibrary Loan Program

It is now 2 years since the new interlibrary loan policy of the National Library of Medicine went into effect. During fiscal year 1959 the National Library of Medicine filled over 6000 interlibrary loan requests each month, and there are many indications that the volume of business will continue in a steady rise.

Two-thirds of all interlibrary loan requests received at the National Library are now being completed within 5 working days. A major objective of the program has been attained in that the percentage of publications "unavailable by reason of being already on loan" has dropped from 33 percent in 1957 to 18 percent in 1958 and to 12 percent in 1959.

The proportion of interlibrary loans being furnished to foreign libraries (one request out of every seven) has remained about the same this year as last. The number of such loans amounted to 10,000 in 1959, leading to speculation as to whether the various national facilities are being fully exploited.

Copies of a revision of the National Library of Medicine's interlibrary loan rules may be obtained on request.

Women in the Class of 1957

For the third successive year, the Women's Bureau of the Department of Labor, with the cooperation of the National Vocational Guidance Association, has conducted a survey of first jobs of women college graduates—nearly 88,000 of them for June 1957. A small percentage had gone on to graduate work, either full or part time—principally in education. Teaching also was the most popular profession of the working graduates, 42,000 being employed in that profession.

In the social, physical, and biological sciences and in mathematics and engineering, however, where many excellent opportunities exist for college women, the numbers were disappointing. Only 586 women had jobs as chemists and 703 as mathematicians and statisticians, or 1.8 percent of employed graduates. "Technicians, biological," represented 2.8 percent.

Since first jobs were largely in fields related to undergraduate majors, it becomes obvious that women are still not aware of the expanding opportunities in the scientific professions. This report by Jean A. Wells of the Women's Bureau emphasizes the importance of proper counseling if young women are to fulfill their individual goals and be of maximum service to society.

A comparison of salaries by occupation reveals that girls who went to work as chemists were the elite of the group, starting at an average annual salary of \$4847. Mathematicians and statisticians did almost as well at \$4675, and biological technicians commanded an average annual salary of \$3854. The over-all annual average was \$3739 (an increase of \$598 since 1955, the first year in which a survey was made). Women who entered journalism were almost as badly paid as typists and clerical workers.

Crerar Library Merger Announced

Merger of the John Crerar Library, one of the world's largest collections of scientific and technical literature, with the Illinois Technological Institute has been announced. The Crerar Library will be housed in a \$2-million public library building to be constructed on the institute's campus.

As a result of this consolidation, the Crerar Library will have a staff of 85 people. With increased facilities and personnel, it is planning improved and expanded services and hopes to realize


substantial economies in library operations. At the same time, the merger will permit the institute to provide increased library services and reference sources for its students, faculty members, and scientific personnel, who comprise at this time Crerar's largest single group of users.

The library will remain under the administration of the Crerar board of trustees and will continue to serve the greater Chicago area in its present capacity. Custody of Illinois Tech's estimated 125,000 scientific and engineering books and periodicals will be assigned to the Crerar Library. The library, with its current inventory of some 1 million books and periodicals of all classifications, plus this acquisition, will have one of the most comprehensive collections of scientific and technological documentation in the United States.

New Helicopter Flight Simulator

A helicopter flight simulator has been developed jointly by the Franklin Institute and the Bell Helicopter Corp. of Philadelphia, Pa. It was built primarily for human engineering studies required in the Army-Navy Instrumentation Program. Because of the problems associated with flight safety, weather, and weight limitations of the helicopter, observation of human performance in actual machines is difficult.

The new simulator can reproduce the motion, sound, vibration, and general cockpit arrangement of most existing helicopters. The pilot operates the test machine as though he were flying a real ship under blind-flight conditions, so that new controls and instruments can be evaluated before being included in a research plane.



The new helicopter flight simulator designed by the Franklin Institute of Philadelphia is used by engineers of the Bell Helicopter Corporation's Army-Navy Instrument Program to evaluate new controls and instruments before they are installed in research helicopters.

Scientists in the News

Donald R. Chadwick, career medical officer of the Public Health Service, has been appointed secretary of the new Federal Radiation Council. Chadwick's appointment was recommended by George B. Kistiakowsky, Special Assistant to the President for Science and Technology and adviser to the Federal Radiation Council.

Pearce Bailey, director of the National Institute of Neurological Diseases and Blindness, National Institutes of Health, Bethesda, Md., has been appointed director of the institute's new International Neurological Research Programs. **Richard L. Masland**, assistant director, will succeed Bailey as director of the institute.

Hilliard Roderick, nuclear physicist in the Controlled Thermonuclear Branch of the Research Division of the U.S. Atomic Energy Commission, has been appointed deputy director of the Natural Sciences Department of the United Nations Educational, Scientific, and Cultural Organization's secretariat in Paris. He started his 2-year assignment in September.

Herbert G. Andrewartha of the University of Adelaide, Adelaide, Australia, has been appointed visiting professor of zoology at Duke University for the fall semester. He will deliver a series of 30 lectures on the physiological aspects of animal distribution.

Sidney H. Liebson, manager of the research and development operations of the Nuclear Development Corporation of America, has been appointed assistant director of physics research at the Armour Research Foundation of Illinois Institute of Technology.

Chester E. Poetsch, head of the pharmaceutical chemistry section of Smith, Kline and French, has been appointed research director of Vick divisions, Vick Chemical Company, New York. He will be vice president of research for Vick Products Division and for Vick International Division.

William C. Coombs, communication engineer, has been appointed chief of the systems analysis section in the Radio Communication and Systems Division of the National Bureau of Standards' Boulder Laboratories, Boulder, Colo.

William E. Adams, James Nelson and Anna Louise Raymond professor of surgery at the University of Chicago, has been appointed chairman of the department of surgery.

Henry P. Wheeler, Jr., chief of the Helium Liaison Office of the U.S. Department of the Interior, has been appointed assistant director of the department's Bureau of Mines, to administer helium activities. He succeeds **Clifford W. Seibel**, who retired on 31 August.

Also at the Department of the Interior, **Paul V. Mullins**, chief of helium operations, has been appointed general manager of helium operations, with headquarters at Amarillo, Tex.

George L. Curran, associate professor of medicine and gerontology at the University of Kansas Medical School, has been appointed associate professor of internal medicine at the St. Louis University School of Medicine. He will direct the section of preventive medicine and public health of the department of internal medicine.

Lee A. DuBridge, president of California Institute of Technology, delivered four lectures, entitled "An Introduction to Space," at Brookhaven National Laboratory, the first in the honorary Pegram lectureship series. The lectureship has been established in the name of the late George B. Pegram, outstanding nuclear physicist and dean of Columbia University, who played an important role in the founding of Brookhaven.

Morris Cohen, professor of physical metallurgy at Massachusetts Institute of Technology, will receive the Francis J. Clamer Medal of the Franklin Institute, Philadelphia, Pa., on 21 October.

Sidney Golden, professor of chemistry at Brandeis University, will spend the coming year at Cambridge University, in the department of theoretical chemistry.

W. A. D. Anderson, professor and chairman of the department of pathology at the University of Miami School of Medicine, Miami, Fla., has received the annual award of the Scientific Products Foundation for the most outstanding contribution to the improvement of laboratory techniques. The presentation took place at the annual meeting of the College of American Pathologists, in Chicago.

Arnold W. Frutkin, secretary to the International Relations Committee of the Space Science Board of the National Academy of Sciences, has been named director of the National Aeronautics and Space Administration's Office of International Programs. He succeeds **Henry E. Billingsley**.

Michael J. S. Dewar, professor and head of the department of chemistry at Queen Mary College, University of London, England, has been appointed professor of chemistry at the University of Chicago.

Charles D. Coryell, professor of chemistry at Massachusetts Institute of Technology and cofounder of the Nuclear Science and Engineering Corp., Pittsburgh, Pa., is the recipient of the American Chemical Society's 1960 Award for Nuclear Applications in Chemistry.

Jean F. Piccard, professor emeritus of aeronautical engineering of the University of Minnesota, has been honored by the American Meteorological Society for his work in the development of balloons for atmospheric research.

Recent Deaths

Sir Ian Heilbron, London, England; 72; specialist on the chemistry of vitamins A and D and on penicillin; had held chairs of organic chemistry at Liverpool University, Manchester University, and the Imperial College of Science and Technology, London; 14 Sept.

Walter B. Kiener, Lincoln, Neb.; 64; botanist and lichenologist; formerly biologist for the Nebraska State Game, Forestation, and Parks Commission; 24 Aug.

Frederick S. Mallette, Riverside, Conn.; 53; specialist on air-pollution control; executive secretary of the Committee on Air-Pollution Controls of the American Society of Mechanical Engineers; 10 Sept.

Arthur Pap, New Haven, Conn.; 37; associate professor of philosophy at Yale University; 7 Sept.

William C. Stadie, Narberth, Pa.; 73; specialist on diabetes; John Herr Musser emeritus professor of research medicine of the University of Pennsylvania; developed an arterial puncture for the study of respiration and cardiac disease and one of the first oxygen tents for the treatment of pneumonia; 11 Sept.

Book Reviews

Asa Gray. A. Hunter Dupree. Belknap Press of Harvard University Press, Cambridge, Mass., 1959. 506 pp. Illus. \$7.50.

Almost every botanist in the United States has, as a student, used Asa Gray's *Manual of Botany*, which is still one of the more useful aids for studying the flora of our northeastern states, and still a very useful tool for teaching students how to identify plants. To beginning botanists, "Gray" is practically synonymous with *Manual of Botany*. Biologists who are interested in evolution (and in 1959, the 100th anniversary of the publication of Darwin's *Origin of Species*, nearly all biologists are) know that when Darwin and Wallace presented their famous papers to the Linnaean Society of London, Darwin included in his, as evidence for his priority in describing natural selection, a letter he had written to Asa Gray. Today, we remember Gray both as a leading systematic botanist of the 19th century and as a personal friend of Charles Darwin—a friend who aided Darwin in establishing and spreading the theory of evolution. How much more there was to Asa Gray, what kind of a man he really was, and what he accomplished as a scientist is told in this remarkably well-written biography by Hunter Dupree.

Now the life of a respectable 19th-century botanist, who labored over his *desiccata* ("baled hay" to irreverent scoffers), who collected specimens over a wide territory, who assembled a herbarium, and who named new genera and species, may be deadly dull even though the subject of the biography may have engaged in the no-holds-barred and no-quarter-given fights over nomenclature. To both students and teachers, reading the biographies of such worthy men may be an unpleasant but necessary chore. But reading this life of Asa Gray is no chore at all. It is a pleasure; in fact, the author has made his book so interesting that most readers will resent, rather than welcome, any little incident that interrupts them. *Asa Gray* is an excellent example of the

kind of writing that has recently made biographies so popular.

This is not intended to imply that Dupree has fictionalized his subject. On the contrary, he has documented it thoroughly by citing over 800 references to primary sources, and if he peers into the subject's mind at times and tells us exactly what Gray's thoughts were, he is always able to cite Gray himself as the authority. In this biography Gray definitely comes alive. But, perhaps, despite his narrative skill and mastery of his subject we need not give the author all of the credit for the interest that this book arouses, for Asa Gray cooperated with his biographer by leading a very interesting life.

Gray was born in 1810 in Sauquoit, New York, and grew up in the central valley, at just the time that the first educational institutions of the region were being established. He became a doctor of medicine in 1831, but practiced only a little more than a year. His primary interests were always those of a naturalist, and he spent all the time that he could spare in gathering botanical, zoological, and mineral specimens. He had the good luck to meet and fall under the influence of John Torrey and, in working with Torrey, he became an expert taxonomist. After some years of odd jobs and financial insecurity, but years of productive scientific labors, he established an outstanding reputation as a botanist, and in 1842 was offered an endowed chair at Harvard University, where his salary of \$1000 a year relieved his financial insecurity. But, to earn his stipend, he had to become a one-man department of botany. He taught botany, established what later became the Gray Herbarium, and supervised the Harvard Botanic Garden. But over and beyond these routine occupations, he indulged his wider interests and spent as much time as he could in identifying and classifying the western plants sent him by various governmental exploring parties.

As Gray's knowledge of the American flora increased, he discovered regularities in the distribution of certain

genera and species, and his interest in plant distribution grew. In fact, he was one of the founders of plant geography, and it was in this field that his very original work came to the attention of Charles Darwin. It was here that, without knowing it, he made an important contribution to the theory of evolution. Gray and Darwin exchanged letters and became close, personal friends, and when Darwin published *The Origin of Species*, Gray reviewed it and saw to it that Darwin got a fair hearing in the United States. Gray was Darwin's first American sponsor.

From a personal standpoint, Gray's long and sometimes bitter contest with his Harvard colleague, Louis Agassiz (who did not accept Darwinian evolution) furnishes the material for perhaps the most interesting portion of the book. Here, we get a glimpse of academic politics, of the growth of science in the first American university, and of the establishment of the National Academy of Sciences. We also get a view of contemporary scientific standards.

Attention should be called to two aspects of this biography that are not adequately expressed in its brief title, *Asa Gray*. First, Dupree's treatment transcends Gray as an individual and depicts, in fascinating detail, the alarms and excursions of 19th-century biology. Dupree makes the science of the time very real, and the actors—the scientists—very human. Finally, he gives us a great many important facts about the group of men who worked with and around Charles Darwin—Sir Charles Lyell, Thomas Henry Huxley, Joseph Hooker, Alfred Russell Wallace and others—the group that introduced evolution to the world. Gray was a very important member of this group.

This biography of Asa Gray is definitive.

CONWAY ZIRKLE

Botanical Laboratory,
University of Pennsylvania

Law and Administration. vols. 1 and 2. Herbert S. Marks, Ed. Pergamon Press, New York, 1959. xiii + 994 pp. \$26.50.

This work is an authoritative and comprehensive source of background information in the field of atomic energy. The tenth of 12 subject categories in the ambitious series "Progress in Nuclear Energy," the work is in two

volumes. Volume 1 presents authoritative articles on specific aspects of law and administration relating to the nuclear-energy programs of the United States, the United Kingdom, Euratom, the International Atomic Energy Agency, and Israel. Volume 2 presents the texts of nuclear-energy legislation and regulations in effect in 25 countries, together with succinct commentaries on the history and status of atomic-energy control in each country.

Each chapter in volume 1 is written by a recognized authority. These chapters present a distillation of the history, conflicts, compromises, and status of the new legal and administrative arrangements that have evolved, primarily in the United States and Britain, as governments respond to the new nuclear technology. John Palfrey's opening chapter is an excellent "legal chronicle" of the United States federal statute, while William Krebs' chapter on state activities and Casper Ooms' chapter on patents are clearly indicative of the authors' deep understanding of the subject matter. I wish that chapters had been included on the new administrative arrangements evolving out of AEC-contractor relationships, or that there had been more specific material concerning the growing federal-state-local problem of licensing and control of private atomic-energy facilities. These are minor criticisms, however, of a noteworthy treatise.

Volume 2, except for the commentaries, is a reference source on foreign legislation and procedures—clearly of limited interest to all but the specialist. The commentaries are useful summaries.

JOHN R. MENKE

*Nuclear Development Corporation
of America, White Plains, New York*

The Emergence of the German Dye Industry. John Joseph Beer. University of Illinois Press, Urbana, 1959. vii + 168 pp. Cloth, \$4.50; paper, \$3.50.

This detailed account of the German dye industry begins with Perkin's discovery of mauve dye (aniline purple) and terminates with the formation of the I. G. Farbenindustrie. It is less a history of the scientific and technical development of the industry than an analysis of the social, economic, and scientific factors which brought about that most

spectacular chapter in the extraordinary history of the chemical industry in the 19th century.

It is surprising that this story has not previously been recounted in English, especially since the German dye industry has been celebrated in the United States as an example of commercial and scientific iniquity. It is the more surprising inasmuch as a considerable number of general histories, company histories, and personal memoirs on the subject in German have long existed.

From this account, the *bête noire* appears to be little worse or better than the rest of us. In fact, while the influence of the chemical industry on German technical education and the patent system was instrumental in securing the initial advantage Germany enjoyed in World War I, Beer concludes that the failure of the German chemical industry consciously to prepare for war in 1914 constituted a limitation on that initial advantage. His conclusions are drawn from published data, plus a few unpublished German sources.

The extent to which Beer brings the German dye industry into the realm of the familiar is illustrated by his remark that "the only reason all the dye companies continued to spend heavily on dye research was for prestige reasons very similar to those which impel the American auto industry to change models every year" (page 123). He emphasizes the role of capitalistic enterprise as a factor in the spectacular growth of the dye companies and notes that Carl Duisberg (later board chairman of I. G. Farbenindustrie) credits to his observations of the American trusts in 1903 some of the ideas which led to the formation of the German combine. The detailed history of the I. G. Farbenindustrie is not a part of this book, however.

Having reduced the story of the German dye industry from the mysterious to the familiar, Beer points out that the factors cited point to the inevitable emergence of the institutional research activities represented by that industry. But he also points out that the coincidences of time and circumstance revealed by an analysis of a specific example such as this are often of the greatest importance in the world as it is. The truth of this is quite evident in the fascinating case history that he has written.

ROBERT P. MÜLTHAUF

Smithsonian Institution

Family Planning, Sterility, and Population Growth. Ronald Freedman, Pascal K. Whelpton, and Arthur A. Campbell. McGraw-Hill, New York, 1959. xi + 515 pp. Illus. \$9.50.

A quarter of a century ago, voluntary parenthood seemed inconsistent with population maintenance. Demographers forecast declining populations with reasonable surety, for birth rates had been moving downward for more than a century. Then birth rates increased in practically all social and economic groups except the lowest, where the decline continued. Moreover, the increases were generally greatest in the groups where the birth rates had been lowest. Voluntary parenthood was consistent not only with childlessness and disappearance of ethnic differences but with a familistic orientation of life that would sustain continuing increase of the population. The assessment of the future population of the nation became a topic of wide conjecture.

The studies made by P. K. Whelpton of the Scripps Foundation for Research in Population Problems had shown that the measurement of trends in fertility should be based on the life-time performance of cohorts of women rather than on the levels of general or age-specific birth rates in specific time periods. However, the past childbearing record of cohorts of women did not provide an empirical basis for estimating future childbearing. These future estimates required knowledge of the wishes and anticipations of families, and of the probable relationship between the numbers of children that were expected and the numbers that would be born. It is these new dimensions of population research that were the goals of the study reported in *Family Planning, Sterility, and Population Growth*.

The Survey Research Center of the University of Michigan and the Scripps Foundation for Research in Population Problems cooperated in this study of family planning and fertility. The 2713 white, married women who were interviewed represented a probability sample of the national population of some 17 million white wives aged 18 to 39 in 1955. The trained interviewers of the Survey Research Center met with unexpected frankness and cooperation in the responses to queries concerning attitudes toward, and practices of, family limitation. Interviews were secured

with 91 percent of the women in the sample and less than 1/2 percent of those interviewed were unwilling to answer questions about contraception.

Summary that does not risk over-generalization is difficult, for the questionnaire was elaborate and the classifications were intricate. There are analyses of sterility and fecundity, the use of contraceptives, and attitudes toward family limitation, methods of limitation, and expected family size; each of these categories is analyzed in relation to the others and to the actual and the desired numbers of children. In addition, all variables are related to such factors as religion, education, income, occupational status, the work of wives, and community background, and again they are considered separately and in relation to each other. Throughout the findings are striking, and they are consistent.

The ghost of biological interpretations of declining fertility is banished by the results of this study. There were no evidences of consistent differences in fecundity (the ability to bear children) between women of the various social and economic strata. The influence of impairments on fertility (actual childbearing) was limited by the fact that couples desired only a few children anyway. However, there are surprising findings in this report. Nine percent of all the women interviewed had been sterilized, many of them for contraceptive purposes. One in four of the wives who had ever been pregnant had had at least one fetal death. And approximately 13 percent of all completed pregnancies had been terminated with a fetal death rather than a live birth.

The fundamental finding of this study of a group of white, married women who constituted a probability sample of the national population was that most couples had a fairly specific idea of how many children they wanted, used contraceptives to space their children and to prevent conception when they had the desired number, and were reasonably successful in limiting the number of children if not always in precise spacing of the children. There were differences among subgroups in the dynamics of family formation, in attitudes toward and use of contraceptives, and in size of the completed family. Religion and educational level were more closely related to fertility variables than were income, occupational status, or community back-

ground, but a majority of the couples in all groups used some control method, and the predominant family in all groups was small. There was nationwide convergence toward the two-, three-, or four-child family as the ideal, the expected, and the fact.

Forecasts of future births utilized the past experience of cohorts of women, the marriage rates and the birth rates of ever-married women in the various cohorts, and the expectations expressed by wives in the study about the size of their completed family. In specific terms:

"The medium fertility projections for ever-married women assume a rise in family size to about 3 births per woman, with a timing pattern similar to that being developed by cohorts of the 1930s. The low series is based on a return to the small families (averaging about 2.5 births) of the cohorts of 1906-15 and a partial return to their older ages at marriage and childbearing. The high series allows for births per ever-married woman to rise to 3.3 (the maximum expectation of the wives interviewed) and for some continuation of the recent tendency to marry and have babies at younger ages" (page 372).

The three series of estimates of births were utilized in combination with conservative assumptions of future declines in mortality and continued net immigration in preparing three series of projections about future populations. The population used as a base was 171 million for the year 1955. The populations as of 1980 were 215 million for the low projection, 239 million for the medium projection, and 262 million for the high projection. The medium projection is the most interesting model, for it assumes a continuation of the marriage and birth-timing patterns of the women born in the 1930's with a completed family size of three births per ever-married woman. This is a small family pattern, and it yields an annual rate of population growth of only 1.3 percent a year in the last quarter of the century. Yet the continuation of the three-child family under the assumed conditions of early marriage and early childbearing within marriage would produce a population of 312 million by the year 2000 and a population of 600 million by the year 2500.

Let me add quickly that the authors stress again and again the implications of their study for continuing and wider

research. The swift development of the two- to four-child family as a nationwide ideal and the almost nationwide achievement of a correspondence between the number of children desired and the actual number of children is incontrovertible evidence that values and behavior in family formation are not static. Repetitive surveys are needed to determine the patterns of correspondence between stated expectations and performance and to ascertain shifts in expectations as they occur. There is the further vast field of research on those factors that are associated with shifts in expectations as they relate to completed fertility, and with the timing of the actual births that lead to a rough equivalence of expectations and completed family size at given levels of expectations.

IRENE B. TAEUBER

*Office of Population Research,
Princeton University*

Ten Steps into Space. Monograph No. 6, *Journal of the Franklin Institute*. Franklin Institute, Philadelphia, Pa., 1958. x + 202 pp. Illus. \$4.

The ten steps of this small paperback book consist of ten lectures (by as many experts) given in a series at the Franklin Institute in the spring of 1958. Although intended for engineers and scientists, most of the material is not highly technical, and much of it is definitely on the popular level. The success of the lecture series led to the decision to attempt to reach a wider audience by publication in book form.

In the first lecture, Ley describes the story of space travel from the earliest speculations to the successes of today, and discusses vividly and simply many of the books, both scientific and science-fiction, that have appeared on the subject. Stehling's lecture, concerned with the fundamental components of rockets and their operation, is carefully developed in nontechnical terms. Richey compares solid, liquid, and nuclear fuels and discusses their characteristics in rather complete technical detail. Singer covers the various kinds of information, such as radiation and magnetic fields, that can be obtained from satellite flights, as well as the experimental techniques involved. Herget succeeds in describing celestial mechanics in a way that involves practically no mathemat-

ics, but he covers the work of Kepler and Newton and the principles of modern, man-made satellite flight. An interesting and detailed description of one particular satellite flight, that of Explorer I, is the subject of Heller's lecture. The most technical account in the book is Ehrcke's detailed mathematical analysis of satellite motion; King's discussion of the known data concerning the atmospheres of Venus and Mars is also technical. Simon gives an interesting description of biological physics, and includes a fascinating description of his own balloon flight to an altitude of approximately 90,000 feet. The series concludes with Levitt's stimulating discussion of the future space possibilities of fission, fusion, and solar energy (solar-sailing and photon-rockets are included); the strange relativistic effects that will occur at speeds approaching the speed of light are also considered.

The eminent scientists who delivered these lectures seem to have made a serious effort to get information across to the nonexpert. In almost all cases, they have succeeded well in accomplishing this challenging task. The level of most of the articles is about right for most of the readers of *Science*, and the book should merit their attention.

DONALD J. HUGHES

Department of Physics,
Brookhaven National Laboratory

A Treasury of Science. Harlow Shapley, Samuel Rapport, and Helen Wright, Eds. Harper, New York, ed. 4, 1958. xiii + 776 pp. Illus. \$6.95.

Anthologies are difficult books to review because the reviewer must exercise a considerable amount of personal and subjective judgment. Either he likes the selections the compilers have included in their book or he doesn't. If the latter, he can always compile his own.

Some years ago I indicated that I liked the *Treasury of Science* [*Science* 104, 430 (1946)]. Two editions later, I still find the book rewarding reading.

Arranged by large topics with numerous subdivisions, the book treats of science and the scientist, the physical world, the world of man, and the rocket and the atom. It is, of course, not possible to list all of the authors included in this scientific smorgasbord. Suffice it to suggest, therefore, that the interested

reader will find tidbits from the writings of Sir J. Arthur Thomson, Patrick Geddes, Albert Einstein, Alan Gregg, and Ivan Pavlov. He will also find selections from Copernicus, Galileo, Newton, Bacon, Darwin, and Huxley.

Although most of the material appeared in the earlier editions, several new sections bring the book closer to today's newspaper headlines. Thus, there is a report on radio astronomy by A. C. B. Lovell, George Wald writes on the origin of life, and Warren Weaver reviews some of the thinking on the genetic dangers of fallout.

In summary, with this new edition *The Treasury* has attained the status of a classic and deserves to be read as well as quoted.

MORRIS C. LEIKIND

National Institute of Neurological
Diseases and Blindness,
National Institutes of Health

Air Pollution Control. W. L. Faith. Wiley, New York; Chapman and Hall, London, 1959. vii + 259 pp. Illus. \$8.50.

This book is intended for those with technical training who are not specialists in the field of air pollution, as well as for intelligent laymen. It is not intended for those seeking a comprehensive treatise on air pollution, for it does not go into any particular phase of air pollution from a fundamental standpoint.

The introductory chapter describes the most important problems of air pollution somewhat superficially. It is surprising that, in the discussion of property damage, there is no mention of the damage to masonry and statuary which occurs in England.

The chapter on meteorology covers general factors that influence the stability of polluted atmospheres. A discussion of smoke problems is followed by a chapter dealing with other particulate materials such as dusts, fumes, and mists. Typical data showing the sources of dustfall and the concentrations encountered in various areas where air pollution exists are given. This material includes a section on sampling and the appraisal of stacks for dusts and fumes as well as brief descriptions of some instruments. In my opinion, sampling procedures should have been treated separately for all cases. Sampling and its concepts assume certain knowledge

which is not readily available to the so-called intelligent layman.

The problem of controlling aerosol emissions is discussed in the section on dust, with a brief résumé of the various devices used, after the discussion of sampling. Again, the treatment is descriptive rather than fundamental.

The discussion of gases is similar to the discussion of the particulate problem, with particular emphasis on sulfur dioxide, hydrogen sulfide, and hydrogen fluoride. Again, sampling methods and control are covered by descriptive materials. Odor problems are discussed in a separate chapter.

Because of the author's personal interest in automobile exhaust, and because of the fact that the Los Angeles area has a serious problem in this respect, the discussion of automobile exhaust and its problems is rather complete. The treatment will be useful to those interested in assessing the present status of the Los Angeles problem.

Separate chapters on air pollution surveys and the legal aspects of air pollution complete the book.

An overall appraisal indicates that the author achieved at least one of his objectives. The book presents some concepts of air pollution to individuals with a technical background. I doubt that it will appeal to the intelligent laymen, as Faith presumed it would.

LESLIE SILVERMAN

School of Public Health,
Harvard University

New Books

The ABC of Relativity. Revised edition edited by Felix Pirani. Allen & Unwin, London; Essential Books, Fair Lawn, N.J., 1959. 139 pp. \$3.75.

Basic Data on Plasma Physics. Sanborn C. Brown. Technology Press and Wiley, New York; Chapman & Hall, London, 1959. 344 pp. \$6.50.

Directory of Nuclear Reactors. vol. 1, *Power Reactors.* International Atomic Energy Agency, Vienna, Austria, 1959 (order from International Publications, 801 Third Ave., New York). 214 pp. Paper, \$3.50.

The Exploration of Space. Arthur C. Clarke. Harper, New York, rev. ed., 1959. 214 pp. \$4.50.

Exploring the Structure of Matter. Jean-Jacques Trillat. Translated by F. W. Kent. Interscience, New York, 1959. 214 pp. \$4.85.

A Field Guide to Bird Songs of Eastern and Central North America. Houghton Mifflin, Boston, Mass., 1959. 2 records, \$10. The records contain the songs and calls of more than 300 species of land and water birds.

Reports

Nitrogen Fixation in Lakes

Abstract. Incorporation of N^{15} into the fixed nitrogen fraction of natural lake waters has been studied for the purpose of estimating rates of primary nitrogen fixation. Experiments in Pymatuning Reservoir, Pennsylvania, in Lake Mendota, Wisconsin, and in two lakes in Alaska indicate that fixation occurs at measurable, sometimes high, rates.

Our understanding of the nature of soil fertility and the availability of nitrogen under natural conditions expanded greatly upon the discovery and subsequent investigation of symbiotic nitrogen fixation in leguminous plants and nonsymbiotic fixation by soil microorganisms. Whether biological fixation of nitrogen *in situ* is a quantitatively important feature of the nitrogen metabolism also of bodies of water is not yet known, although marine and freshwater biologists have sometimes tacitly assumed that this is the case (1). The list of organisms known to be capable of fixing free nitrogen is long; included on it are species of blue-green algae (particularly of the genera *Nostoc* and *Anabaena*, but also of *Tolypothrix*, *Calothrix*, and others), common and often abundant constituents of freshwater plankton, as well as *Azotobacter* and photosynthetic bacteria, including some that have been isolated from the water and mud of Lake Mendota, Wisconsin (2). Organisms capable of fixing nitrogen are therefore known to occur in the aquatic environment; it remains only to show that they con-

tribute significantly to the supply of fixed nitrogen there.

We have used uptake of the stable isotope N^{15} as a measure of nitrogen fixation in lake water. Our method employs preliminary removal of dissolved atmospheric nitrogen from water samples contained in special 1000-ml closed vessels by aeration with an 80:20 helium-oxygen mixture at a pressure of 0.8 atm. Sufficient N_2 containing 95 atom percent N^{15} is then added to bring the pressure within the vessel to approximately 1 atm, and the system is equilibrated by shaking. Samples are incubated either in the lake at the depth from which the water was originally collected or under specified laboratory conditions. Subsequent treatment is essentially the same as that applied by Rittenberg, Keston, Roseburg, and Schoenheimer (3) and by Burris, Eppling, Wahlin, and Wilson (4) in their studies (5).

Preliminary measurements were made on the following lakes in 1958 (samples were incubated in the lake): (i) Little Kitoi and Upper Jennifer lakes, Afognak Island (Kodiak region), Alaska; (ii) Pymatuning Reservoir (Sanctuary Lake), Pennsylvania; (iii) Lake Mendota, Wisconsin.

The amount of N^{15} incorporated into the fixed-nitrogen fraction was estimated in all of these instances by subtracting the measured atom percent N^{15} of laboratory air from that of the fixed-nitrogen fractions of the samples after incubation. It has become obvious that this method overestimates the excess of N^{15} for two reasons. (i) Air is not an adequate standard because it gives a value about 0.002 atom percent lower than N_2 from ammonium sulfate which has gone through the conversion procedure. The latter, therefore, provides a standard more comparable to the actual samples. (ii) Lake-water blanks sterilized and carried through the entire experimental procedure appear to be enriched by about 0.004 atom percent N^{15} , as compared with ammonium sulfate. We have, therefore, applied a correction of -0.006 atom percent to these data, not having used adequate

blanks at the time the measurements were made. Rates computed in this way are obviously not particularly reliable but may be used as preliminary estimates. Seven measurements made in the Alaskan lakes during July and August yielded rates of fixation ranging from 0 to 0.0047 μg of nitrogen per liter per day. Fixation was not detectable in a single experimental series from Lake Mendota in late October. A single series from Pymatuning Reservoir on 4 October yielded rates of incorporation far too high to be canceled by the most conservative corrections for the blank, of the order of 0.8 μg of nitrogen per liter per day. This figure is based on duplicate samples, the results of which agreed within 1 percent. There is no reason to doubt that here nitrogen was fixed at a high rate. For all experiments in 1958, incubation periods ranged from 4 to 6 days.

In more recent experimental series from Pymatuning Reservoir (beginning 20 April 1959) use has been made of certain technical refinements: (i) reagent-grade ammonium sulfate (instead of laboratory air) was used as a primary standard for isotope ratio; (ii) lake-water blanks were carried through the entire procedure after initial autoclaving within the experimental vessels; and (iii) all experimental vessels were autoclaved prior to the introduction of the sample, sterile procedure being observed subsequently. Water for these experiments was collected at Pymatuning, transported to Pittsburgh, and treated as described above about 8 hours after collection; samples were incubated under constant illumination at 15°C (approximately the temperature of the water when it was collected) for varying lengths of time. The results are given in Fig. 1 as atom percent N^{15} (excess over ammonium sulfate and sterile lake-water blank) plotted against




Fig. 1. Incorporation of N^{15} by surface water taken from Sanctuary Lake, Pymatuning Reservoir, Pennsylvania, on 20 Apr. 1959, incubated under constant illumination at 15°C.

Instructions for preparing reports. Begin the report with an abstract of from 45 to 55 words. The abstract should not repeat phrases employed in the title. It should work with the title to give the reader a summary of the results presented in the report proper.

Type manuscripts double-spaced and submit one ribbon copy and one carbon copy.

Limit the report proper to the equivalent of 1200 words. This space includes that occupied by illustrative material as well as by the references and notes.

Limit illustrative material to one 2-column figure (that is, a figure whose width equals two columns of text) or to one 2-column table or to two 1-column illustrations, which may consist of two figures or two tables or one of each.

For further details see "Suggestions to Contributors" [Science 125, 16 (1957)].

time of incubation. The curve appears to be characterized by two rates of fixation, one of about 0.021 μ g of nitrogen per liter per day during the first 58 hours, the other of about 0.068 μ g of nitrogen per liter per day during the rest of the experiment (6).

RICHARD DUGDALE
VERA DUGDALE

Department of Biological Sciences,
Pymatuning Laboratory of Field
Biology, University of Pittsburgh,
Pittsburgh, Pennsylvania

JOHN NEESS
JOHN GOERING

Department of Zoology,
University of Wisconsin, Madison

References and Notes

1. C. E. ZoBell, *Marine Microbiology* (Chronica Botanica, Waltham, Mass., 1946); F. Rutner, *Fundamentals of Limnology* (Univ. of Toronto Press, Toronto, 1953).
2. E. S. Lindstrom et al., *J. Bacteriol.* 61, 481 (1951).
3. D. Rittenberg, A. S. Keston, F. Rosebury, R. Schoenheimer, *J. Biol. Chem.* 127, 285, 291 (1939).
4. R. H. Burris, F. J. Eppling, H. B. Wahlin, P. W. Wilson, *Soil Sci. Soc. Am. Proc.* 7, 258 (1942).
5. A detailed description of our entire procedure is in preparation.
6. We wish to express our appreciation to R. H. Burris of the department of biochemistry, University of Wisconsin, and Richard Abrams of the research laboratory, Montefiore Hospital, Pittsburgh, for the advice they have generously given and for the use of the mass spectrometers in their laboratories. Both spectrometers are Consolidated-Nier isotope ratio machines. This research was supported by the University of Kentucky Faculty Research Fund and by the National Institutes of Health. The Alaska Department of Fish and Game provided laboratory facilities at the Kitoi Bay Research Station on Afognak Island.

13 July 1959

Sensory Deprivation and Visual Speed: An Analysis

Abstract. Speeds of moving objects were markedly underestimated by human observers after prolonged patternless visual stimulation. Even greater underestimation followed exposure to a "noisy" visual field; on the other hand, exposure to a hyperstable field caused overestimation. The effects of external visual noise simulate those of deprivation; this finding suggests that similarly disordered but spontaneous neural discharge dominates the visual nervous system in deprivation.

Experiments in sensory deprivation have been made to explore the consequences of prolonged exposure to visual fields in which normal spatiotemporal variation is sharply reduced. Results of experiments performed at McGill University (1) suggest that the apparent speed of a moving line markedly changes as one consequence of long-term exposure to an illuminated but

patternless visual field. After such deprivation, observers reported that a straight black line slowly rotating against a dimly lit screen looked S-shaped. This distortion was attributed to an induced "perceptual lag": the ends of the line lag behind the center.

In preliminary investigations we found no evidence for a perceived distortion of the line. Instead, the apparent speed of the entire line was clearly reduced. Consequently, in further investigations we have used an apparatus (Fig. 1) designed to measure and thus quantify changes in apparent speed (2). The observer was instructed to fixate a black dot 1 m distant at eye level. He then saw a black bar (Fig. 1, A) sweep, like the second hand of a clock, through 90 deg from a horizontal to a vertical position, at a rate of 60 deg/sec. At the vertical, the bar disappeared from the observer's sight behind a screen. His task was to judge when the now-hidden bar would reach a fixed marker (Fig. 1, B) 10 deg beyond. This judgment was based on the observer's estimate of the speed of the bar during its visible 90 deg of travel. When he judged that the hidden bar had traversed the 10 deg to reach the marker, he stopped the coupled time clock (Fig. 1) which gave a measure of the apparent speed of the bar. The average of the initial judgments made by practiced observers was very close to the actual travel time. Twenty trials per observer were made before and after each experimental exposure.

In a first test to evaluate our interpretation of the McGill results, seven observers were deprived of patterned vision for 8 hours under conditions similar to those of the McGill study (3). The observers, lying on cots, were subjected to a masking acoustic noise at the maximum tolerable level; they wore tubes over their arms and hands and translucent goggles over their eyes. The goggles provided only a homogeneously illuminated, nonpatterned field of vision. After deprivation, each of the seven observers judged the speed of the sweep to be less (by an average of 16 percent) than he had initially judged it to be. The observers showed no accompanying change in simple motor reaction time. Having demonstrated and measured this consequence of generalized deprivation, we proceeded to further analyses.

Each of 12 observers was instructed to fix his gaze on the center of a patternless but illuminated circular field (intensity 0.2 ml) subtending 43 visual degrees in an otherwise dark field (patternless exposure field). Although only visual input was controlled, a significant reduction in apparent speed was pro-

Table 1. Changes in apparent speed after ½ hour of exposure. Exposure fields: H, hyperstable; D, dark; P, patternless; N, noisy.

Exposure field	Change in speed (%)	Significance* of differences between conditions
H	+14.8	.05
D	-4.8	
P	-10.1	N.S.†
N	-19.6	
		.01
		.02
		.001

* Significance levels derived from an analysis of variance. † Not significant.

duced with only ½ hour of exposure (Table 1). This test demonstrates that control of extravisual stimulation is not essential for obtaining the change in apparent speed.

Comparable experimental situations were then used to explore the effects of exposing the same observers to three other test fields devised on the basis of the following speculation. In the absence of normal stimulation coming from patterned fields, there is reason to suspect that the visual nervous system exhibits spontaneous and patternless activity (4). Dominance of the visual system for long periods by such intrinsic noise may result in reduction of




Fig. 1. Apparatus for measuring apparent speed. A, Rotating sweep; B, fixed marker.

apparent speed. According to this reasoning, even greater reduction should result from exposing observers to a barrage of randomized visual stimulation (extrinsic noise). An electronic device was used to produce a continually changing display of dots on the face of a 14-inch television tube. The resulting pattern may be called visual noise and defined as a changing pattern of dots devoid of redundancies of position or intensity (noisy exposure field). At the opposite extreme from visual noise there is stimulation with minimal temporal variation—a condition that can be produced by exposure to a fixed pattern, called here a hyperstable field because of the absence of even that amount of object-movement found in everyday scenes. To achieve comparability of exposure conditions we used a still photograph of the picture-tube display which showed the set of dots available at one instant (hyperstable exposure field). Such stimulation may suppress intrinsic noise to a degree greater than normal and hence cause an increase in apparent speed.

Finally, the observers were kept alert in total darkness for a comparable period (dark exposure field). Under the latter three test conditions, illumination level and field size (except for the dark exposure field), exposure time, and other controlled variables were equated with those of the first test. The order of presentation was varied for the 12 observers. The resulting changes in apparent speed are shown in Table 1. An analysis of variance showed that the differences among conditions were significant beyond the .001 level.

Thus, a method has been developed for quantifying one consequence of visual deprivation. Tests in which this method was used have yielded results consistent with the following propositions: (i) Simulation and enhancement of an aftereffect of visual deprivation through exposure to a noisy visual field implies that deprivation entails the randomization of sensory-neural activity rather than the diminution or absence of such activity. (ii) Production of an aftereffect opposite to that of deprivation, after exposure to a hyperstable field, implies that typical exposure fields have relevant noise characteristics somewhere between those of the noise used and those of the hyperstable field. (iii) The normal stability of speed perception depends upon continuous exposure to the typical environment.

RICHARD HELD
BURTON WHITE

Department of Psychology, Brandeis University, Waltham, Massachusetts

References and Notes

1. W. Heron, B. K. Doane, T. H. Scott, *Can. J. Psychol.* **10**, 13 (1956).
2. This research was supported by a grant from the National Science Foundation.
3. These tests were made in collaboration with S. J. Freedman, director of isolation studies, Massachusetts Mental Health Center.
4. R. Granit, *Receptors and Sensory Perception* (Yale Univ. Press, New Haven, Conn., 1955).
- 11 June 1959

Protection by Sulfur Compounds against the Air Pollutants Ozone and Nitrogen Dioxide

Abstract. Two distinct but related pathways of protection against the lethal effects of ozone and nitrogen dioxide are shown by (i) simultaneous inhalation of compounds that furnish $-SH$ or $-SS-$, or both, and (ii) by injection of thiourea derivatives several days prior to exposure to these oxidant gases. The mechanism of (i) is believed similar to that proposed for the action of radiation-protective compounds; that of (ii) involves the development of a tolerance initiated by the thiourea against the oxidants.

Ozone (O_3) and nitrogen dioxide (NO_2) are potent respiratory irritants which may be injurious when inhaled (1, 2), and evidence exists that the concentration of O_3 in urban air is damaging to plants (3). Nitrogen dioxide, in addition to acting as a pulmonary irritant, acts as a precursor of O_3 in oxidant smogs by photochemical dissociation (4).

In tests designed to determine the acute toxicologic effects of combinations of either known or suspected air pollutants, we found that certain sulfur compounds were effective in counteracting the toxic action of O_3 and NO_2 . Partial protection by ascorbic acid against acute lethal effects of O_3 has been previously reported from our lab-

oratories (5) as well as by Mittler (6), who also found slight protection by sodium thiosulfate. Mittler's conclusion that "compounds which protect against death by radiation . . . are not effective against ozone poisoning" is not supported by our work, however. Tests involved the inhalation exposure of mice to an approximate LC_{50} of either O_3 or NO_2 simultaneously with one or more sulfur compounds. In some instances the sulfur compounds were administered intraperitoneally prior to inhalation of the oxidant. Mortality resulting from a 4-hour exposure, as compared with the mortality from a control exposure to oxidant alone, reflected enhancement or suppression of O_3 or NO_2 toxicity as influenced by the sulfur compound. Methods of generation and analysis of O_3 and vapor concentrations of sulfur compounds have been described elsewhere (5, 7). Gaseous sulfur compounds and NO_2 were similarly administered from cylinders in metered amounts and diluted to the desired concentration with purified air. The Saltzman method was used for NO_2 analysis (8).

The data in Table 1 represent selected results of a series of tests which show the maximal protection found for each sulfur compound. Comparison of mortalities in sulfur-treated versus control groups indicates the degree of protection afforded. Hydrogen sulfide (H_2S) gave significantly greater protection against oxidant exposure, particularly NO_2 , than other sulfur compounds. On a molar basis H_2S protected in a ratio of 1/55 moles of NO_2 , whereas benzenethiol, next in order of effectiveness, protected in a ratio of 1/5.4. Higher molar ratios of sulfur compounds were required for protection against O_3 , however; for benzenethiol 1.5 mole/mole of O_3 , and for H_2S , 2 mole/mole of O_3 .

Table 1. Effect of sulfur compounds on the toxicity of nitrogen dioxide and ozone for mice. Mortality at 24 and 72 hours is indicated by deaths/number of mice tested.

Sulfur compound	Av. concn. (ppm)	Av. concn. of oxidant (ppm)	Mortality			
			S treated		Oxidant alone	
			24 hr	72 hr	24 hr (†)	72 hr
1-Hexanethiol	145	78	0/20	1/20	10/20	11/20
1-Hexanethiol	115	4.1	1/20	2/20	10/20	10/20
Methanethiol	65	4.8	2/15	2/15	9/15	9/15
Dimethyl disulfide	45	83	5/20	5/20	10/20	11/20
Dimethyl disulfide	21 mg/kg*	80	4/20	4/20	13/20	13/20
Dimethyl disulfide	21 mg/kg*	4.6	2/20	2/20	10/20	12/20
Hydrogen polysulfide	20 mg/kg*	105	8/25	8/25	16/25	18/25
Di-tert-butyl disulfide	24	84	4/15		9/15	
Benzenethiol	14	76	1/20	1/20	10/20	10/20
Benzenethiol	9	6.1	1/20	3/20	11/20	12/20
Hydrogen sulfide†	11	4.9	7/35	7/35	17/35	18/35
Hydrogen sulfide†	1.5	82	1/20	2/20	10/20	10/20
Thiophene	180	85	9/15		8/15	
Dimethyl sulfide	195	4.6	14/30		14/30	

* Administered by intraperitoneal injection. † An aged technical grade.

Certain —SS— compounds also gave protection against oxidant toxicity; both dimethyl disulfide and hydrogen polysulfide significantly protected intraperitoneally injected mice, and dimethyl disulfide and *tert*-butyl disulfide protected via inhalation. It is noted that purified H₂S provided less protection than that shown in Table 1 for a technical grade. This difference in protective ability is attributed on infrared evidence to the presence of —SS—, possibly hydrogen polysulfide, in the technical H₂S, and inasmuch as addition of —SS— to purified H₂S reinforced its protective ability.

The antagonism of oxidant toxicity by sulfur compounds is physiologic and not the result of chemical combination prior to inhalation to form less active compounds. Analyses of chamber atmosphere for oxidant gas showed essentially no change upon addition of sulfur compounds. Moreover, protection was conferred by the injected compounds. Histologic examination of pulmonary tissue showed degrees of tissue change that paralleled the protective effect. Pulmonary edema and cellular infiltration, characteristic responses of oxidant exposure, were markedly inhibited in the protected groups.

The mechanisms involved in the protection are, as yet, unknown. The functional unit appears to be —SH or —SS—, or both, but not —S—; dimethyl-sulfide and thiophene were ineffective. Significantly, —SH and —SS— are characteristic of compounds conferring protection against ionizing radiation (9). Brinkman and Lamberts (10) have called attention to the possible radiomimetic properties of O₃ by showing that O₃ and irradiation produced the same defect in oxygen consumption of the skin of the finger; also cysteamine was equally protective against the effects of both agents. Likewise, Fetner (11) has shown a similar capacity of O₃ and x-rays to produce chromosomal aberrations in *Vicia faba*. The action was presumably mediated through the OH and HO₂ radicals formed from the aqueous decomposition of O₃; the separate effects of O₃ plus irradiation were fully additive.

Single injections of MEG (2-mercaptoethylguanidine-HBr), highly protective against radiation at 200 mg/kg, gave modest protection against NO₂ (mortality: 8/20 S treated, versus 12/20 for NO₂ alone) and O₃ (13/20 versus 19/20). When given in a series of three injections (100 mg/kg on each of 3 days), however, MEG provided greater protection (2/20 versus 10/20 for NO₂ alone). Similarly, 0.04 mg/kg of BAL (2,3-dimercaptopropanol) provided some protection to mice exposed to NO₂ (6/22 versus 12/22 for NO₂ alone) but did not decrease O₃ toxicity (13/25 versus 12/25 for O₃ alone).

Thus, there is similarity of action of protective compounds which favors the view that the mechanism of action of the oxidants O₃ and NO₂ may be in part similar to that of x-irradiation.

Another means of protecting against the lethal effects of NO₂ and O₃ was achieved by still other sulfur compounds, α -naphthylthiourea (ANTU) and phenylthiourea (PTU). Mice exposed to 5.8 ppm of O₃ the day following the last of 3 intraperitoneal injections of ANTU (10 mg/kg every other day) and rats which received 128 ppm of NO₂ 18 days after administration of ANTU (8 mg/kg) showed marked tolerance to the oxidants. Comparative mortalities were: mice, 0/15 versus 12/15 for controls; rats, 1/6 versus 6/6 for controls. Similarly, PTU (8 mg/kg) given in four injections every other day and exposed to either O₃ or NO₂ 3 days after the last injection produced good protection (4/10 versus 9/10 for O₃ alone, and 4/10 versus 8/10 for NO₂ alone). Although the mechanism presumably still involves reactive sulfur constituents, the pathway is obviously different from that mediated by —SH and —SS— compounds because the tolerance afforded by ANTU and PTU persists, a condition not met by the simple —SH and —SS— compounds. Also, unlike —SH and —SS—, simultaneous treatment with oxidant and ANTU (itself a potent producer of pulmonary edema) produces additive toxicity, instead of protection. It has been demonstrated that rats develop a marked tolerance to ANTU (12) and also to O₃ (1). The cross tolerance between ANTU and oxidant indicates that the basic mechanisms may be similar. Furthermore, it appears that the tolerance mechanism is related to the edema mechanism, and the latter to sulfur balance, inasmuch as —SH (cysteine) blocks the lethal effects of ANTU (13), whereas —SH or —SS—, or both, reduces oxidant toxicity.

It is felt that the antagonism to oxidants displayed by certain sulfur compounds is highly significant for the insight it provides concerning possible mechanisms for protection against air pollutants, and also toward elucidation of the hitherto unexplained toxic action of oxidants (14).

EDWARD J. FAIRCHILD II
SHELDON D. MURPHY
HERBERT E. STOKINGER
*Occupational Health Program, U.S.
Public Health Service, Cincinnati, Ohio*

References and Notes

1. H. E. Stokinger, *A.M.A. Arch. Ind. Health* **14**, 181 (1957).
2. W. F. von Oettingen, *U.S. Public Health Service, Public Health Bull. No. 272*, Security Agency, Public Health Service (1941).
3. H. T. Freebairn, *Science* **126**, 303 (1957).
4. A. J. Haagen-Smit, *Proc. 2nd Natl. Air Pollution Symposium, Pasadena, Calif.* (1952).

5. R. N. Matzen, *J. Appl. Physiol.* **11**, 105 (1957).
6. S. Mittler, *Nature* **181**, 1064 (1958).
7. E. J. Fairchild and H. E. Stokinger, *Am. Ind. Hyg. Assoc. J.* **19**, 171 (1958).
8. B. E. Saltzman, *Anal. Chem.* **26**, 1949 (1954).
9. D. G. Doherty, W. T. Burnett, Jr., R. Shapira, *Radiation Research* **7**, 13 (1957); R. Shapira, D. G. Doherty, W. T. Burnett, Jr., *ibid.* **7**, 22 (1957); A. Pihl and L. Eldjarn, *Pharmacol. Revs.* **10**, 437 (1958).
10. R. Brinkman and H. B. Lamberts, *Nature* **181**, 1202 (1958).
11. R. W. Fetner, *ibid.* **181**, 504 (1958).
12. C. P. Richter, *J. Am. Med. Assoc.* **129**, 927 (1945).
13. K. P. DuBois, L. W. Holm, W. L. Doyle, *Proc. Soc. Exptl. Biol. Med.* **61**, 102 (1946).
14. We thank L. D. Scheel of our laboratory for preparing the hydrogen polysulfide, and K. P. DuBois, professor of pharmacology, University of Chicago, for supplying the ANTU.

16 July 1959

Uridine Isomer (5-Ribosyluracil) in Human Urine

Abstract. A substance isolated from human urine by anion exchange absorption and paper chromatography was found to correspond in its ultraviolet absorption spectrum and chromatographic mobilities with a uridine isomer (5-ribosyluracil) recently described as a component of yeast ribonucleic acid.

In 1957 Davis and Allen (1) reported the isolation from yeast ribonucleic acid of a new nucleotide ("the fifth nucleotide") and described its physical and chemical properties, notably a characteristic bathochromic shift in the ultraviolet absorption spectrum at alkaline pH. The corresponding nucleoside, prepared by the action of intestinal phosphatase, was subjected to hydrazinolysis, and the carbohydrate component was identified as D-ribose. Cohn (2) also obtained from yeast ribonucleic acid an "apparently modified uridylic acid" identical with the "fifth nucleotide" of Davis and Allen.

The structure of this new nucleotide proved to be of unusual interest. Recent investigation by Yu and Allen (3) revealed that the nitrogenous component is uracil and that the D-ribose, which has the furanose configuration, apparently is attached to the pyrimidine ring at the 5 position. Methylation studies by Scannell, Crestfield, and Allen (4) also indicated the structure to be 5-ribosyluracil. Further evidence for this formulation was obtained by Cohn (5) by means of periodate oxidation and nuclear magnetic resonance spectra. The C—C ribosyl linkage reported for the uridine isomer is unique thus far for components of nucleic acid, but C—C glycosyls occur in certain natural products (6).

A compound identical in spectral and chromatographic properties with this isomer of uridine was encountered in

Table 1. Spectral data for the urinary substance and for 5-ribosyluracil from yeast ribonucleic acid (from 3).

pH	$\lambda_{\text{max.}}$ (m μ)	$\lambda_{\text{min.}}$ (m μ)	Ratios			
			250/260	280/260	290/260	300/260
Yeast Ribonucleic Acid						
2	263	234	0.74	0.43	0.07	
12	286	245	0.64	2.09	2.16	1.22
Urine						
2	263	233	0.75	0.43	0.08	
12	287	246	0.65	2.09	2.19	1.29

the course of an investigation of nucleosides occurring in urine in man (7). The urine of two adult males was studied. One was a normal subject; the other was gouty. Both abstained from consumption of tea and coffee for the period of study. The urine specimens were adjusted to pH 9 with ammonium hydroxide and were filtered. The filtrate was allowed to percolate through an amount of Dowex 2 (8 \times , 20-50 mesh) ion-exchange resin in acetate form equivalent to one-sixth of its volume. After washing-in with ammonium acetate buffer (pH 9) and water, the column was eluted with water. The aqueous eluate was collected in $\frac{1}{2}$ column-volume fractions until the initial $E_{260\text{m}\mu}^{1\text{cm}}$ of 15.0 decreased to a value of 0.3. The combined fractions were evaporated to dryness under reduced pressure and at temperatures below 50°C. The residue was extracted with 60 percent aqueous ethanol. The extract brought to dryness *in vacuo* was dissolved in a minimum amount of water and applied to Whatman 3 MM filter-paper sheets for descending two-dimensional chromatography (8). A solvent system composed of *n*-butyl alcohol and 0.6M ammonium hydroxide (6:1) (9) was used in the long dimension of the sheet for 58 hours, and a system composed of *n*-butyl alcohol, formic acid, and water (77:11:12) (10) in the second direction for 17 hours. A spot which strongly

absorbed ultraviolet light, located nearest the origin of the chromatogram, contained the uridine isomer. An extract of this spot gave the absorption spectrum shown in Fig. 1. The close correspondence of the spectral data with those reported for the uridine isomer derived from yeast ribonucleic acid by Yu and Allen (3) is indicated in Table 1.

A comparison of the chromatographic mobilities of the two compounds in four solvent systems was made (11). The relative mobility values of the spot extract coincided with those of authentic 5-ribosyluracil. Thus, in solvent system A [isobutyric acid, ammonia (0.5) (10:6)], the relative mobilities of the authentic sample and of the urinary product were 0.88 and 0.85, respectively (mobility of uridine = 1.0); in solvent system B [isopropanol, acetic acid, water (6:3:1)], 0.66 and 0.60; in solvent system C [tertiary amyl alcohol, water, formic acid (6:3:1)], 0.76 and 0.73; in solvent system D [butanol, water (86:14)], 0.53 and 0.48.

With three solvents (A, B, and C) the urinary product appeared homogeneous. The fourth solvent (D), however, revealed a small residual spot at its origin, indicating some impurity. We therefore used solvent D for further purification of a larger sample by band chromatography, obtaining a microcrystalline product.

A rough estimate of the amounts of the uridine isomer excreted in 24 hours was obtained by applying the figure for $\epsilon_{\text{max}} = 7.5 \times 10^3$ at pH 2 given by Yu and Allen (3) to the spot extracts. Values of 18 mg/24 hr for the urine of the normal subject and 41 mg/24 hr for the urine of the gouty subject were obtained. Further studies are in progress to determine whether these differences in the gouty and nongouty subjects are consistent. In any event, both figures are in excess of the quantities of free purine bases reported previously in human urine (8).

The possibility that the uridine isomer found in the urine is derived from the diet is not excluded, although foods rich in nucleic acids, as well as coffee and tea, were omitted by the two sub-

jects investigated. An endogenous origin is considered more likely, however. This would imply that the new uridine isomer occurs in human tissue, and this is in accord with the reported presence of the isomer in dog pancreas (12). It may belong to the class of naturally occurring "additional" nitrogenous constituents of ribonucleic acid which thus far are known to include 5-methylcytosine, various methylated purines, and thymine (13).

MAX ADLER

ALEXANDER B. GUTMAN

Mount Sinai Hospital and
College of Physicians and Surgeons,
Columbia University, New York

References and Notes

1. F. F. Davis and F. W. Allen, *J. Biol. Chem.* **227**, 907 (1957).
2. W. E. Cohn, *Federation Proc.* **16**, 166 (1957); **17**, 203 (1958).
3. C. Yu and F. W. Allen, *Biochim. et Biophys. Acta* **32**, 393 (1959).
4. J. P. Scannell, A. M. Crestfield, F. W. Allen, *ibid.* **32**, 406 (1959).
5. W. E. Cohn, *ibid.* **32**, 569 (1959).
6. E. J. Hay and L. J. Haynes, *J. Chem. Soc.* **1956**, 3141 (1956); **1958**, 2231 (1958); W. H. Evans, A. McGookin, J. Jurd, E. Robertson, W. R. N. Williamson, *ibid.* **1957**, 3510 (1957).
7. This study was supported in part by grant A-162 from the National Institute of Arthritis and Metabolic Diseases, U.S. Public Health Service.
8. B. Weissmann, P. A. Bromberg, A. B. Gutman, *J. Biol. Chem.* **224**, 407, 423 (1957).
9. E. Chargaff, R. Lipshitz, C. Green, M. E. Hodes, *ibid.* **192**, 223 (1951).
10. R. Markham and J. D. Smith, *Biochem. J.* **45**, 294 (1949).
11. We are indebted to Frank W. Allen (see 1) for chromatographic identification.
12. J. W. Kemp and F. W. Allen, *Biochim. et Biophys. Acta* **28**, 51 (1958).
13. M. Adler, B. Weissmann, A. B. Gutman, *J. Biol. Chem.* **230**, 717 (1958); J. W. Littlefield and D. B. Dunn, *Biochem. J.* **70**, 642 (1958); H. Amos and M. Korn, *Biochim. et Biophys. Acta* **29**, 444 (1958).

15 June 1959

Production of Large Amounts of Plant Tissue by Submerged Culture

Abstract. The growth rate of plant tissue cultures is substantially increased through the use of a large-volume carboy system. Aeration is considered to be the most important factor. With this system, yields of a pound or more of tissue can be obtained within 2 weeks.

The methods of plant tissue culture have suggested a way to put the metabolic systems of plants to work under controlled conditions. A prerequisite for the most efficient use of this idea would be the large-scale production under submerged conditions of the tissue or tissues involved. Although the growth of microorganisms under these conditions is now commonplace, and advances in this direction with animal cells and tissues have been made (1), no such work has been reported for plants.




Fig. 1 Absorption spectra of the urinary uridine isomer.




Fig. 1. The carboy culture apparatus. Aeration and agitation are accomplished by the introduction of compressed air (on the left) through a side-arm trap and then through two Seitz filters, which sterilize the air.

Since this approach constitutes part of the tissue-culture program in this laboratory, some of our results on the large-volume growth of plant tissue cultures are presented here.

The principal tissue used for large-volume culture was one obtained from Paul's Scarlet rose stems. This tissue was selected because of its rapid growth rate on solid media and the fact that it grows well in liquid culture. The medium used for carboy studies was made up of White's basal medium (2), 2, 4-dichlorophenoxyacetic acid (6 parts per million), yeast extract (0.1 percent), and malt extract (0.1 percent).

The inoculum for the carboy is prepared by taking the tissue culture from a test tube and placing it in liquid medium in a 300-ml erlenmeyer flask. This culture is then agitated on a New Brunswick gyrorotatory shaker. After 1 week's




Fig. 2. Growth curves for Paul's Scarlet rose stem tissue grown in carboy culture and in erlenmeyer flasks. The growth value equals the ratio (final fresh weight: initial fresh weight).

growth, the contents of one flask (the culture in 100 ml of medium) is transferred to a Fernbach flask containing 900 ml of medium. This flask is also aerated and agitated by shaking. After an additional week's growth, the Fernbach culture (1000 ml) is inoculated into a 20-liter carboy containing 9 liters of medium. This is the final stage of this particular scale-up. For purposes of aeration and agitation of the culture during growth, the carboy is fitted with stainless-steel tubing to receive compressed air, which is sterilized by passage through two Seitz filters (Fig. 1). In addition, appropriate connections are included for an air outlet, the addition of medium, and sampling. In all of these manipulations, aseptic procedures must be followed.

Samples of the carboy culture were made at weekly intervals and compared with results obtained from the same tissue grown on a New Brunswick gyrorotatory shaker in 300-ml erlenmeyer flasks containing 100 ml of culture. These results, shown in Fig. 2, indicate an accelerated rate of growth in the carboy for the first 2 weeks. Growth values (final wet wt./initial wet wt.) of 5.5 and 14.9 are attained by the tissue grown in the carboy system after 1 week and 2 weeks, respectively, as compared to growth values of 1.1 and 1.9 for the same periods with the shaker cultures.

The difference in growth rates between these two systems during the first week is due to the method of aeration. At the end of the first week of growth, 2 liters of medium were added to the carboy; the effect of this supplementation is an increase in rate of growth during the second week. When no further additions are made, the rate of growth drops off sharply, as shown by the growth values for the third and fourth weeks.

Other tissues also grow well in this carboy system of culture. The data available thus far for tissues from *Ilex* stem, *Lolium* endosperm (3), and *Ginkgo* pollen (4) indicate growth increases similar to the increase found with the Paul's Scarlet rose stem. The *Ginkgo* pollen tissue, for example, yielded 2 lb of cells after 4 weeks in carboy culture.

Other factors recognized as important in increasing the growth rate of plant tissue cultures are those well known to microbiologists working with large-volume cultures. Besides the necessity for an adequate nutrient medium, which itself may be improved in numerous ways, a number of physical factors are involved. In addition to the method and efficiency of aeration, which have already been stressed, the temperature of incubation and the means of agitation are important. More dependent on the tissue culture itself are such

things as the age of the inoculum and the particular strain of tissue selected for growth. Moreover, the maintenance of a maximum growth rate rests on such things as timely addition of medium supplements, minimizing of foam production, and proper buffering of the hydrogen ion concentration.

From a practical standpoint, the carboy system of culturing plant tissues is a step in the direction of decreasing the lag phase of growth and increasing the ultimate yield. Our results emphasize the importance of timely nutritional supplements to the medium and of adequate aeration for the rapid growth of certain plant tissue cultures. The aeration factor receives major consideration in microbial "fermentations," but investigations with plant material have not been concerned with the growth of large amounts of material. In the roller-tube technique employed by Steward, Caplin, and Millar (5) and the variable-speed shaker technique used by Muir, Hildebrandt, and Riker (6), aeration has been studied, but the aim has not been the production of large yields (7). Another difference is that in both of these systems the agitation provides the aeration, whereas in the carboy system the aeration provides the agitation and at the same time furnishes a continuous flow of air.

WALTER TULECKE
LOUIS G. NICKELL

Phytochemistry Laboratory,
Chas. Pfizer and Company,
Brooklyn, New York

References

1. W. F. McLimans et al., *J. Immunol.* **79**, 428 (1957); *J. Bacteriol.* **74**, 768 (1957).
2. P. R. White, *Handbook of Plant Tissue Culture* (Ronald, New York, 1943).
3. K. J. Norstog, *Botan. Gaz.* **117**, 253 (1956).
4. W. Tulecke, *Am. J. Botany* **44**, 602 (1957).
5. F. C. Steward, S. M. Caplin, F. K. Millar, *Ann. Botany (London)* **16**, 57 (1952).
6. W. H. Muir, A. C. Hildebrandt, A. J. Riker, *Am. J. Botany* **45**, 589 (1958).
7. L. G. Nickell, *Proc. Natl. Acad. Sci. U.S.* **42**, 848 (1956).

6 June 1959

Pipecolic Acid in Leaves of Strawberry Plant as Influenced by Treatments Affecting Growth

Abstract. Growth inhibition due to maleic hydrazide spray or to an unsatisfied chilling requirement caused young fully expanded leaves of the strawberry to accumulate pipecolic acid to a very striking degree. It is postulated that the accumulation resulted from blockage in normal metabolic conversion to other intermediary nitrogenous compounds.

Pipecolic acid is a naturally occurring cyclic imino acid that has been isolated from several plants and identified chromatographically in many others (1, 2).

Table 1. Nitrogenous fractions in young fully expanded strawberry leaves, in relation to treatments with maleic hydrazide (MH), 1957.

Treatment and date	Nitrogenous fractions		
	Total* ($\mu\text{g N/g}$ fresh wt.)	Pipecolic acid $\mu\text{g N/g}$ fresh wt.	% of total N
<i>Variety: Sparkle</i>			
None			
8/22	60.1	0.4	0.7
9/12	73.3	1.3	1.9
MH, 1000 ppm			
8/22	56.6	0.4	0.8
9/12	74.3	4.8	6.5
MH, 2000 ppm			
8/22	54.0	0.5	1.0
9/12	95.1	6.6	7.0
<i>Variety: Catskill</i>			
None			
8/25	78.9	1.0	1.4
9/12	81.3	1.1	0.4
MH, 1000 ppm			
8/25	65.3	1.4	2.3
9/12	94.6	3.2	3.4
MH, 2000 ppm			
8/25	73.1	1.6	2.2
9/12	78.4	9.5	9.7

* Alcohol-soluble ninhydrin-reacting substances.

Usually present in low concentration, it has been found to accumulate strikingly in the immature fruit and embryo of *Phaseolus vulgaris* (2). Working with that species, Grobbelaar and Steward (3) have found that lysine is converted to pipecolic acid, which in turn may be metabolized to α -amino adipic acid.

In the strawberry plant (*Fragaria chiloensis* var. *ananassa*) pipecolic acid is usually not found in very significant concentration among the alcohol-soluble ninhydrin-reacting nitrogenous compounds that are separated by two-directional paper chromatography. However, recent studies have indicated that this compound may accumulate dramatically in leaves of strawberry plants which are internally inhibited from active growth in spite of exposure to favorable growing conditions.

In the first of these studies, field-grown plants of the varieties Sparkle and Catskill were sprayed with aqueous solutions containing 1000 and 2000 parts of maleic hydrazide (MH) per million on 20 August and again on 4 September 1957. By 6 September, the growth-inhibiting effects of the MH sprays were visible, but there was no apparent injury to the leaf surface. Between 20 August and 15 September samples of young fully expanded leaves were taken at intervals of 2 to 3 days and preserved in 80-percent alcohol. The alcohol extracts were subsequently analyzed for amino acids by two-directional chromatography, with phenol

and butanol-acetic acid used as the solvents. The pipecolic acid spots were identified by position and by their fluorescence in ultraviolet light. Quantitative estimation was accomplished by the procedure of Thompson and Morris (4). Table 1 shows the pipecolic acid concentrations (micrograms of N per gram fresh weight) in young leaves from untreated and MH-treated plants early and late in the experimental period. The rise in pipecolic acid due to MH was apparent in the Sparkle leaf samples by approximately 12 days after the first treatment, and accumulation continued progressively until the last sampling time.

In the second study, young Sparkle strawberry daughter plants were potted in late August and held outside until early November. They were then divided into two lots, one of which was held in the greenhouse at 65° to 75°F under 10-hour days, and the other subjected to chilling at 35° to 45°F continuously, also under short-day conditions. Approximately 2 months later the chilled plants were brought to the greenhouse. At that time both the chilled plants and those that had been held continuously at warm temperatures were separated into long (16-hour) day and short (12-hour) day lots. After a month, these four lots of plants were again divided, and half of each lot was sprayed with 100 parts of gibberellic acid (Gibrel) per million.

The unchilled plants grew more slowly and their leaves were smaller and darker green than those from chilled plants in all lots. In addition to the growth stimulus from chilling, there were additive stimulatory effects of long days and Gibrel. Thus, the unchilled, short-day plants that received no Gibrel grew least, and the chilled, long-day plants that received Gibrel grew most. Table 2 shows the effects of these treatments on pipecolic acid content, as micrograms per gram of dry weight, of the young fully expanded leaves sampled about 2 weeks and about 5 weeks after the chilled plants were brought to the greenhouse. At the first sampling the growth stimulus resulting from chilling was accompanied by a very great rise in ninhydrin-reacting nitrogenous compounds. As a result, although no consistent difference in the micrograms of N as pipecolic acid was apparent, that present amounted to 13.8 and 14.9 percent of the total soluble nitrogen accounted for in leaves of unchilled plants and only 3.6 and 4.9 percent of that found in leaves from chilled plants. Three weeks later, and about 2 weeks after Gibrel treatment, ninhydrin-reacting N had increased markedly in all the leaf samples from nonchilled plants and had decreased in the leaves from chilled

Table 2. Influence of chilling, day length, and gibberellic acid treatment on nitrogenous fractions in young, fully expanded leaves of the Sparkle strawberry, 1959.

Date and treatment	Nitrogenous fractions		
	Total* ($\mu\text{g N/g}$ dry wt.)	Pipecolic acid $\mu\text{g N/g}$ dry wt.	% of total N
<i>Short day</i>			
(1/30)			
Chilled	1283	46	3.6
Nonchilled	398	59	14.9
(2/19)			
Chilled	654	16	2.4
Nonchilled	769	146	18.9
(2/19)			
Chilled + Gibrel	733	14	1.9
Nonchilled + Gibrel	833	193	23.2
<i>Long day</i>			
(1/30)			
Chilled	1091	54	4.9
Nonchilled	279	39	13.8
(2/19)			
Chilled	485	11	2.4
Nonchilled	757	300	39.6
(2/19)			
Chilled + Gibrel	529	14	2.7
Nonchilled + Gibrel	570	96	16.8

* Alcohol-soluble ninhydrin-reacting substances.

plants. In addition, there was a striking increase in pipecolic acid N in all four leaf samples from unchilled plants, and this became the dominant form of ninhydrin-reacting N in those samples, rising to 39.6 percent of the total in the nonchilled plants that were under long days but did not receive Gibrel.

Lysine was low in all samples and did not vary consistently as a result of treatments affecting growth. If α -amino adipic acid was present, it did not appear at the expected position on any of the two directional chromatograms in sufficient concentration to react with ninhydrin. It seems possible, therefore, that the increase in pipecolic acid associated with internal growth inhibition resulted from blockage in metabolic conversion to other compounds. (5).

LAWRENCE YATSU
DAMON BOYNTON

Department of Pomology,
Cornell University, Ithaca, New York

References and Notes

1. F. C. Steward and J. K. Pollard, *Ann. Rev. Plant Physiol.* 8, 65 (1957).
2. N. Grobbelaar, doctoral thesis, Cornell Univ. N.Y. (1955).
3. — and F. C. Steward, *J. Am. Chem. Soc.* 75, 4341 (1953).
4. J. F. Thompson and C. J. Morris, *Anal. Chem.*, in press.
5. We are grateful to J. F. Thompson and J. K. Pollard for advice and material assistance. This investigation was supported in part by research grant G5588 from the National Science Foundation.

30 April 1959

THE AMERICAN ASSOCIATION FOR THE ADVANCEMENT OF SCIENCE and THE NATIONAL SCIENCE FOUNDATION

announce the publication of revised editions of two popular booklists, and the issuance of three new publications of the Science Library Program:

THE TRAVELING HIGH SCHOOL SCIENCE LIBRARY (5th edition, 1959), an annotated catalog of the 200 books in a special loan collection being sent to 1700 schools this year, 25¢ a copy.

THE TRAVELING ELEMENTARY SCHOOL SCIENCE LIBRARY (1st edition), an annotated catalog of the 160 books in a special collection being sent to 800 selected elementary schools, 25¢ a copy.

AN INEXPENSIVE SCIENCE LIBRARY (3rd edition, 1959), a selected and annotated list of over 400 paperbound science books suitable for high-school students and non-specialist adult readers, 25¢ a copy.

THE AAAS SCIENCE BOOK LIST (1959), an annotated checklist of about 900 books recommended for high-school and public libraries prepared with the collaboration of members of the AAAS Council and other scientists and educators, \$1.00 per copy, net.

A SELECTED LIST OF CAREER GUIDANCE PUBLICATIONS, prepared for high school students and teachers. Single copy free.

A discount of one-third is offered on orders for 25 or more copies of all lists, *except The AAAS Science Book List*, if sent to a single address. Remittances should accompany orders totalling less than \$5.00. *Do not send postage stamps in payment.*

Address orders to AAAS PUBLICATIONS
1515 Massachusetts Avenue, N.W.
Washington 5, D.C.

Meetings

Forthcoming Events

November

1-4. Society of Economic Geologists, Pittsburgh, Pa. (H. M. Bannerman, U.S. Geological Survey, Washington 25.)

2-4. American Clinical and Climatological Assoc., Hot Springs, Va. (F. T. Billings, 420 Medical Arts Bldg., Nashville, Tenn.)

2-4. Atomic Industrial Forum, annual conf., Washington, D.C. (Atomic Industrial Forum, Inc., 260 Madison Ave., New York 16.)

2-4. Geochemical Soc., Pittsburgh, Pa. (K. B. Krauskopf, Geology Dept., Stanford Univ., Stanford, Calif.)

2-4. Geological Soc. of America, Pittsburgh, Pa. (H. R. Aldrich, 419 W. 117 St., New York 27.)

2-4. Mineralogical Soc. of America, Pittsburgh, Pa. (C. S. Hurlbut, Jr., 12 Geological Museum, Harvard Univ., Oxford St., Cambridge 38, Mass.)

2-4. National Assoc. of Geology Teachers, Pittsburgh, Pa. (F. Foote, Dept. of Geology, Williams College, Williamstown, Mass.)

2-4. Paleontological Soc., Pittsburgh, Pa. (H. B. Whittington, Museum of Comparative Zoology, Harvard Univ., Cambridge 38, Mass.)

2-5. Physical and Extractive Metallurgy, symp., Chicago, Ill. (Metallurgical Soc. of AIME, 29 W. 39 St., New York 18.)

2-6. American Inst. of Mining, Metallurgical, and Petroleum Engineers and Inst. of Metals, fall, Chicago, Ill. (E. O. Kirkendall, AIME, 29 W. 39 St., New York 18.)

2-6. Collegium Internationale Allegalogicum, 4th symp., Rome, Italy. (A. Cerletti, Pharmacological Laboratories, Sandoz Ltd., Basel, Switzerland.)

4-5. Diffraction, 17th annual conf., Pittsburgh, Pa. (P. K. Koh, Allegheny Ludlum Steel Corp., Research and Development Laboratories, Brackenridge, Pa.)

4-6. American Nuclear Soc., conf., Washington, D.C. (American Nuclear Soc., John Crerar Library, 86 E. Randolph St., Chicago 1, Ill.)

4-6. Antibiotics, 7th annual symp., Washington, D.C. (H. Welch, Div. of Antibiotics, Food and Drug Administration, Dept. of Health, Education, and Welfare, Washington 25.)

4-6. Design of Experiments in Army Research, 5th conf. (by invitation only), Fort Detrick, Frederick, Md. (F. G. Dressel, Office of Ordnance Research, Box CM, Duke Station, Durham, N.C.)

4-6. Eastern Analytical Symp., New York, N.Y. (P. Lublin, Publicity Chairman, Sylvania Research Laboratories, Bay-side, N.Y.)

4-6. Industrial management Soc., Chicago, Ill. (R. J. Mayer, IMS, 330 S. Wells St., Chicago 6.)

4-6. National Automatic Control Conf., Dallas, Tex. (G. L. Turin, Hughes Research Laboratories, Culver City, Calif.)

4-6. Society of Rheology, 30th anniversary, Bethlehem, Pa. (J. T. Bergen, Armstrong Cork Co., Lancaster, Pa.)


4-6. Technical Assoc. of the Pulp and

A black and white photograph of a collection of geometric objects, including rectangular blocks, cylinders, and a tall, thin structure, arranged on a dark surface. The objects are illuminated from the side, creating strong highlights and shadows.

Klett-Summerson Photoelectric Colorimeters—
Colorimeters—Nephelometers—Fluorimeters—
Bio-Colorimeters—Comparators—Glass Stand-
ards—Klett Reagents.

LEISS

Single and Double
MIRROR-MONOCHROMATORS



with exchangeable prisms for the visible, ultraviolet, infrared from 200 millimicrons to 20 microns

Write for Bulletin #980 to

PHOTOVOLT CORP.
95 Madison Ave. New York 16, N.Y.

All specific information including a tabulation of principal isotopes now arranged for greater clarity • Nobelium 102 (most recent discovery) included

Chart and 48-page KEY booklet both completely revised by Dr. William F. Meggers, National Bureau of Standards

PERIODIC CHART OF THE ATOMS																		Atomic Number	
The Atoms Grouped According to the Number of Outer (Valence) Electrons																		Planetary electrons in the completed shells	
Atomic Number	Symbol	Symbol	Symbol	Symbol	Symbol	Symbol	Symbol	Symbol	Symbol	Symbol	Symbol	Symbol	Symbol	Symbol	Symbol	Symbol	Symbol	Symbol	
1	H	2	He	3	Li	4	Be	5	B	6	C	7	N	8	O	9	F	10	Ne
11	Na	12	Mg	13	Al	14	Si	15	P	16	S	17	Cl	18	Ar	19	K	20	Ca
21	Sc	22	Ti	23	V	24	Cr	25	Mn	26	Fe	27	Co	28	Ni	29	Cu	30	Zn
31	Ga	32	Ge	33	As	34	Se	35	Br	36	Kr	37	Rb	38	Sr	39	Y	40	Zr
41	Nb	42	Mo	43	Tc	44	Ru	45	Rh	46	Pd	47	Ag	48	Cd	49	In	50	Sn
51	Sb	52	Te	53	I	54	Xe	55	Cs	56	Ba	57	La	58	Ce	59	Pr	60	Nd
61	Pm	62	Sm	63	Eu	64	Gd	65	Tb	66	Dy	67	Ho	68	Er	69	Tm	70	Yb
71	Lu	72	Hf	73	Ta	74	W	75	Re	76	Os	77	Ir	78	Pt	79	Au	80	Hg
81	Tl	82	Pb	83	Bi	84	Po	85	At	86	Rn	87	Fr	88	Ra	89	Ac	90	Th
91	Pa	92	U	93	Np	94	Pu	95	Am	96	Cm	97	Bk	98	Cf	99	Es	100	Fm
101	La	102	Ce	103	Pr	104	Nd	105	Pm	106	Sm	107	Eu	108	Gd	109	Tb	110	Dy
111	Lu	112	Hf	113	Ta	114	W	115	Re	116	Os	117	Ir	118	Pt	119	Au	120	Hg
121	Tl	122	Pb	123	Bi	124	Po	125	At	126	Rn	127	Fr	128	Ra	129	Ac	130	Th
131	Pa	132	U	133	Np	134	Pu	135	Am	136	Cm	137	Bk	138	Cf	139	Es	140	Fm
141	La	142	Ce	143	Pr	144	Nd	145	Pm	146	Sm	147	Eu	148	Gd	149	Tb	150	Dy
151	Lu	152	Hf	153	Ta	154	W	155	Re	156	Os	157	Ir	158	Pt	159	Au	160	Hg
161	Tl	162	Pb	163	Bi	164	Po	165	At	166	Rn	167	Fr	168	Ra	169	Ac	170	Th
171	Pa	172	U	173	Np	174	Pu	175	Am	176	Cm	177	Bk	178	Cf	179	Es	180	Fm
181	La	182	Ce	183	Pr	184	Nd	185	Pm	186	Sm	187	Eu	188	Gd	189	Tb	190	Dy
191	Lu	192	Hf	193	Ta	194	W	195	Re	196	Os	197	Ir	198	Pt	199	Au	200	Hg
201	Tl	202	Pb	203	Bi	204	Po	205	At	206	Rn	207	Fr	208	Ra	209	Ac	210	Th
211	Pa	212	U	213	Np	214	Pu	215	Am	216	Cm	217	Bk	218	Cf	219	Es	220	Fm
221	La	222	Ce	223	Pr	224	Nd	225	Pm	226	Sm	227	Eu	228	Gd	229	Tb	230	Dy
231	Lu	232	Hf	233	Ta	234	W	235	Re	236	Os	237	Ir	238	Pt	239	Au	240	Hg
241	Tl	242	Pb	243	Bi	244	Po	245	At	246	Rn	247	Fr	248	Ra	249	Ac	250	Th
251	Pa	252	U	253	Np	254	Pu	255	Am	256	Cm	257	Bk	258	Cf	259	Es	260	Fm
261	La	262	Ce	263	Pr	264	Nd	265	Pm	266	Sm	267	Eu	268	Gd	269	Tb	270	Dy
271	Lu	272	Hf	273	Ta	274	W	275	Re	276	Os	277	Ir	278	Pt	279	Au	280	Hg
281	Tl	282	Pb	283	Bi	284	Po	285	At	286	Rn	287	Fr	288	Ra	289	Ac	290	Th
291	Pa	292	U	293	Np	294	Pu	295	Am	296	Cm	297	Bk	298	Cf	299	Es	300	Fm
301	La	302	Ce	303	Pr	304	Nd	305	Pm	306	Sm	307	Eu	308	Gd	309	Tb	310	Dy
311	Lu	312	Hf	313	Ta	314	W	315	Re	316	Os	317	Ir	318	Pt	319	Au	320	Hg
321	Tl	322	Pb	323	Bi	324	Po	325	At	326	Rn	327	Fr	328	Ra	329	Ac	330	Th
331	Pa	332	U	333	Np	334	Pu	335	Am	336	Cm	337	Bk	338	Cf	339	Es	340	Fm
341	La	342	Ce	343	Pr	344	Nd	345	Pm	346	Sm	347	Eu	348	Gd	349	Tb	350	Dy
351	Lu	352	Hf	353	Ta	354	W	355	Re	356	Os	357	Ir	358	Pt	359	Au	360	Hg
361	Tl	362	Pb	363	Bi	364	Po	365	At	366	Rn	367	Fr	368	Ra	369	Ac	370	Th
371	Pa	372	U	373	Np	374	Pu	375	Am	376	Cm	377	Bk	378	Cf	379	Es	380	Fm
381	La	382	Ce	383	Pr	384	Nd	385	Pm	386	Sm	387	Eu	388	Gd	389	Tb	390	Dy
391	Lu	392	Hf	393	Ta	394	W	395	Re	396	Os	397	Ir	398	Pt	399	Au	400	Hg
401	Tl	402	Pb	403	Bi	404	Po	405	At	406	Rn	407	Fr	408	Ra	409	Ac	410	Th
411	Pa	412	U	413	Np	414	Pu	415	Am	416	Cm	417	Bk	418	Cf	419	Es	420	Fm
421	La	422	Ce	423	Pr	424	Nd	425	Pm	426	Sm	427	Eu	428	Gd	429	Tb	430	Dy
431	Lu	432	Hf	433	Ta	434	W	435	Re	436	Os	437	Ir	438	Pt	439	Au	440	Hg
441	Tl	442	Pb	443	Bi	444	Po	445	At	446	Rn	447	Fr	448	Ra	449	Ac	450	Th
451	Pa	452	U	453	Np	454	Pu	455	Am	456	Cm	457	Bk	458	Cf	459	Es	460	Fm
461	La	462	Ce	463	Pr	464	Nd	465	Pm	466	Sm	467	Eu	468	Gd	469	Tb	470	Dy
471	Lu	472	Hf	473	Ta	474	W	475	Re	476	Os	477	Ir	478	Pt	479	Au	480	Hg
481	Tl	482	Pb	483	Bi	484	Po	485	At	486	Rn	487	Fr	488	Ra	489	Ac	490	Th
491	Pa	492	U	493	Np	494	Pu	495	Am	496	Cm	497	Bk	498	Cf	499	Es	500	Fm
501	La	502	Ce	503	Pr	504	Nd	505	Pm	506	Sm	507	Eu	508	Gd	509	Tb	510	Dy
511	Lu	512	Hf	513	Ta	514	W	515	Re	516	Os	517	Ir	518	Pt	519	Au	520	Hg
521	Tl	522	Pb	523	Bi	524	Po	525	At	526	Rn	527	Fr	528	Ra	529	Ac	530	Th
531	Pa	532	U	533	Np	534	Pu	535	Am	536	Cm	537	Bk	538	Cf	539	Es	540	Fm
541	La	542	Ce	543	Pr	544	Nd	545	Pm	546	Sm	547	Eu	548	Gd	549	Tb	550	Dy
551	Lu	552	Hf	553	Ta	554	W	555	Re	556	Os	557	Ir	558	Pt	559	Au	560	Hg
561	Tl	562	Pb	563	Bi	564	Po	565	At	566	Rn	567	Fr	568	Ra	569	Ac	570	Th
571	Pa	572	U	573	Np	574	Pu	575	Am	576	Cm	577	Bk	578	Cf	579	Es	580	Fm
581	La	582	Ce	583	Pr	584	Nd	585	Pm	586	Sm	587	Eu	588	Gd	589	Tb	590	Dy
591	Lu	592	Hf	593	Ta	594	W	595	Re	596	Os	597	Ir	598	Pt	599	Au	600	Hg
601	Tl	602	Pb	603	Bi	604	Po	605	At	606	Rn	607	Fr	608	Ra	609	Ac	610	Th
611	Pa	612	U	613	Np	614	Pu	615	Am	616	Cm	617	Bk	618	Cf	619	Es	620	Fm
621	La	622	Ce	623	Pr	624	Nd	625	Pm	626	Sm	627	Eu	628	Gd	629	Tb	630	Dy
631	Lu	632	Hf	633	Ta	634	W	635	Re	636	Os	637	Ir	638	Pt	639	Au	640	Hg
641	Tl	642	Pb	643	Bi	644	Po	645	At	646	Rn	647	Fr	648	Ra	649	Ac	650	Th
651	Pa	652	U	653	Np	654	Pu	655	Am	656	Cm	657	Bk	658	Cf	659	Es	660	Fm
661	La	662	Ce	663	Pr	664	Nd	665	Pm	666	Sm	667	Eu	668	Gd	669	Tb	670	Dy
671	Lu	672	Hf	673	Ta	674	W	675	Re	676	Os	677	Ir	678	Pt	679	Au	680	Hg
681	Tl	682	Pb	683	Bi	684	Po	685	At	686	Rn	687	Fr	688	Ra	689	Ac	690	Th
691	Pa	692	U	693	Np	694	Pu	695	Am	696	Cm	697	Bk	698	Cf	699	Es	700	Fm
701	La	702	Ce	703	Pr	704	Nd	705	Pm	706	Sm	707	Eu	708	Gd	709	Tb	710	Dy
711	Lu	712	Hf	713	Ta	714	W	715	Re	716	Os	717	Ir	718	Pt	719	Au	720	Hg
721	Tl	722	Pb	723	Bi	724	Po	725	At	726	Rn	727	Fr	728	Ra	729	Ac	730	Th
731	Pa	732	U	733	Np	734	P												

W. M. WELCH SCIENTIFIC COMPANY
DIVISION OF W. M. WELCH MANUFACTURING COMPANY
ESTABLISHED 1880
1515 Sedawick Street, Dept. E, Chicago 10, Illinois, U.S.A.

No. 4854

- Improved readability
- Large size—42 X 58 inches
- Lithographed in 6 colors
- Lists latest atomic weights compiled by International Union for Pure & Allied Chemistry

The Welch Periodic Chart of the Atoms is the most authoritative and comprehensive presentation of the principal facts about all known atoms including the artificial transuranic elements recently manufactured. The information is sufficiently detailed to serve the most advanced workers in atomic physics, yet it presents basic facts clearly to students of elementary high school chemistry and physics. Its organization permits constant visualizing of each characteristic in its proper place with respect to the whole picture of atomic structure.

4854. CHART OF THE ATOMS. 1959 Edition.

With formed-metal chart molding at top and bottom with eyelets for hanging, non-glare, protective coating, and Key booklet. **Each, \$7.50**

Paper Industry, 13th alkaline pulping conf., Jacksonville, Fla. (TAPPI, 155 E. 44 St., New York 17.)

5-8. Group for the Advancement of Psychiatry, New York, N.Y. (American Psychiatric Assoc., 1700 18 St., NW, Washington 9.)

6. Gastroenterology Research Group, 9th semi-annual, Chicago, Ill. (E. Clinton Texter, Jr., Ward Memorial Bldg., Medical School, Northwestern Univ., 303 E. Chicago Ave., Chicago 11.)

8-13. International Rubber Conf., Washington, D.C. (B. S. Garvey, Jr., Pennsalt Chemical Corp., Industrial Chemicals Div., 813 Lancaster Pike, Wayne, Pa.)

9-11. American Petroleum Inst., 39th annual, Chicago, Ill. (API, 50 W. 50 St., New York 20.)

9-11. Association of Military Surgeons, 66th annual conv., Washington, D.C. (R. E. Bitner, AMS, Suite 718, 1726 Eye St., NW, Washington 6.)

9-11. Chemical Engineering, symp., Hamilton, Ontario, Canada. (Chemical Inst., 18 Rideau St., Ottawa 2, Ontario.)

9-11. Institute of Radio Engineers—Electronics Industries Assoc., fall, Syracuse, N.Y. (L. G. Cumming, IRE, 1 E. 79 St., New York 21.)

9-11. Instrumentation Conf., 4th, Atlanta, Ga. (W. B. Jones, Jr., School of Electrical Engineering, Georgia Inst. of Technology, Atlanta 13.)

9-12. Society of Exploration Geophysicists, 29th annual intern, Los Angeles, Calif. (B. Roberts, SEG, 1544 N. Highland Ave., Los Angeles 28.)

10-12. Electrical Techniques in Medicine and Biology, 12th annual conf., Philadelphia, Pa. (D. A. Holaday, College of Physicians and Surgeons, Columbia Univ., New York 32.)

10-15. Laboratory Measurement and Automation Techniques in Chemistry, intern. cong., Basel, Switzerland. (ILMAC, 61 Clarastrasse, Basel, Switzerland.)

11-12. Clinical Anticancer Drug Research, Washington, D.C. (B. H. Morrison, III, Cancer Chemotherapy National Service Center, National Cancer Inst., Bethesda 14, Md.)

11-13. Gerontological Soc., Detroit, Mich. (R. W. Kleemeier, Dept. of Psychology, Washington Univ., St. Louis 5.)

11-14. Society of Naval Architects and Marine Engineers, annual, New York, N.Y. (W. N. Landers, SNAME, 74 Trinity Pl., New York 6.)

12-13. Cardiology in Aviation, intern. symp., Brooks Air Force Base, Tex. (L. E. Lamb, Dept. of Internal Medicine, School of Aviation Medicine, Brooks Air Force Base.)

12-13. Operations Research Soc., natl., Pasadena, Calif. (D. A. Katcher, 4608 Morgan Drive, Chevy Chase 15, Md.)

12-13. Utilization of Atomic Energy, 2nd annual conf., College Station, Tex. (G. M. Krise, Radiation Biology Laboratory, Texas Engineering Experiment Station, College Station.)

12-18. International Odontological Session (with 64th Paris Dental Congress), Paris, France. (J. Charon, Secretary-General, 31, rue Tronchet, Paris 8^e, France.)

15-18. Society of American Foresters, 59th, San Francisco, Calif. (Soc. of American Foresters, Mills Bldg., 17th and Pennsylvania Ave., NW, Washington 6.)

15-19. American Soc. of Agronomy, Cincinnati, Ohio. (L. G. Monthey, 2702 Monroe St., Madison 5, Wisc.)

15-20. Radiological Soc. of North America, conf., Chicago, Ill. (Radiological Soc. of North America, 815 Medical Arts Bldg., Fort Worth, Tex.)

16-18. Molecular Structure, 3rd conf., Houston, Tex. (Robert A. Welch Foundation, 2010 Bank of the Southwest Bldg., Houston 2.)

16-19. Magnetism and Magnetic Materials, 5th conf., Detroit, Mich. (D. M. Grimes, Dept. of Electrical Engineering, Univ. of Michigan, Ann Arbor.)


16-20. American Rocket Soc., annual meeting, Washington, D.C. (J. J. Harford, ARS, 500 Fifth Ave., New York 36.)

16-20. Automation Cong., 5th intern., New York, N.Y. (R. Rimbach, 845 Ridge Ave., Pittsburgh 12, Pa.)

16-21. Antarctic Symp., Buenos Aires, Argentina. (R. N. Panzarini, Instituto Antartico, Argentino, Cerrito 148, Buenos Aires.)

16-21. Disposal of Radioactive Waste, conf., Monaco. (Intern. Atomic Energy Agency, 11-13 Kärntner Ring, Vienna 1, Austria.)

17-19. Building Research Inst. (NAS-NRC), fall conf., Washington, D.C. (J. H. Houtchens, Information Services, BRI, NAS-NRC, 1145 19 St., NW, Washington 25.)



74147B

74217E



74217D



LOW COST / ALOE U-V CUVETTES

ROUND / MATCHED / ULTRAVIOLET TRANSMITTING

For use with
Beckman DU Spectrophotometers

Available for the first time, these low cost, Aloe, ultraviolet transmitting, round cuvettes offer distinct advantages to all educational, industrial and other research laboratories. Multiple problem of low cost, accurate U-V cuvettes now solved with these round, matched cuvettes. Proven in laboratory use over a year. No unusual etching noted when 4 normal acid or alkali concentrations were allowed to stand for as long as two hours.

They compare most favorably in accuracy, to say nothing of cost, with conventional square "quartz" cuvettes in the 235-340 mμ range where 95% of all U-V measurements are made.

74147A — Aloe U-V Cuvettes, round, set of 4, matched to tolerance of ½% T.....\$20.00

74147B — Aloe U-V Cuvettes, round, set of 12, matched to tolerance of 1% T.....\$24.00

Accessories:

Precision Molded Adapter — holds cuvettes in firm position.

74217F — Aloe U-V Cuvette Adapter, Each.....\$5.00

74217E — Aloe U-V Cuvette Adapter with matched set of 4 cuvettes.\$25.00

Aloe U-V Cuvette Adapter Carriage — precision made, yet versatile.

74217D — Aloe U-V Cuvette Adapter Carriage, Each.....\$32.00


Direct your order or request for additional information to:

aloe scientific 5655 Kingsbury, St. Louis 12, Mo.

DIVISION OF THE BRUNSWICK-BALKE-COLLENDER COMPANY

FULLY STOCKED DIVISIONS COAST-TO-COAST





Write for this free 40-page book

HARSHAW SCINTILLATION PHOSPHORS


Presents definitive article on characteristics and properties of scintillation phosphors with special emphasis on NaI(Tl). A general discussion of scintillation counting is augmented with many appropriate tables, efficiency curves, and typical gamma ray spectra. Gives specifications and drawings of Harshaw mounted phosphors, and lists miscellaneous other phosphors available from Harshaw. We will be pleased to send you a copy.

Also available . . .

Free 36-page book

HARSHAW SYNTHETIC OPTICAL CRYSTALS

Discusses in detail various Harshaw crystals used for infra-red and ultra-violet optics. Includes many pertinent graphs. Ask for your copy today.




THE HARSHAW CHEMICAL COMPANY

1945 EAST 97th STREET • CLEVELAND 6, OHIO


CHICAGO • CINCINNATI • CLEVELAND • DETROIT • HOUSTON • LOS ANGELES • PHILADELPHIA
HASTINGS-ON-HUDSON, N. Y. • PITTSBURGH

COUNTERCURRENT FRACTIONATOR



↑ **EC520 COUNTERCURRENT FRACTIONATOR** with automatic drive; 2 racks of 100 20-ml tubes on base 24" x 34".

PAPER AND STARCH ELECTROPHORESIS



↑ **EC451 RESEARCH MODEL** with migration chamber 20 x 45cm. Water-cooled pressure plates for paper, sponge, or starch electrophoresis.

SEND FOR FURTHER INFORMATION AND DETAILED SPECIFICATIONS

E-C APPARATUS COMPANY—538 WALNUT LANE—SWARTHMORE, PA.

GET YOUR ADVANCE COPY

of the General Program of the AAAS Chicago Meeting by first class mail – early in December

The General Program of the 126th Meeting of the AAAS in Chicago, 26-31 Dec., 1959, will be available to you within the first week in December—whether you can attend the Meeting or not.

Effective this year, the former General Program-Directory, which had become an unwieldy book of more than 400 pages, has been separated into *two* publications, namely:

- a) The Directory of AAAS Officers and Activities, 96 pp., already published; and
- b) The General Program of the Annual Meeting, c. 200 pp., which will appear early in December

Both of these, sold at cost, may be purchased separately—in advance (see coupon below), or at the meeting. Some of their *respective* contents are:

The General Program

1. The two-session general symposium "Moving Frontiers of Science IV," arranged by the Committee on AAAS Meetings.
2. Programs of the 18 AAAS sections (symposia and contributed papers).
3. Programs of the more than 70 participating societies.
4. Sessions of the Conference on Scientific Communication, Conference on Scientific Manpower, and the Academy Conference.
5. The Special Sessions: AAAS Address and Reception, National Geographic Society, Phi Beta Kappa, Sigma Xi, RESA, Tau Beta Pi Association.
6. Details of the Morrison Hotel—center of the Meeting—and of the other session sites.
7. Titles of the latest foreign and domestic scientific films to be shown in the AAAS Science Theatre.
8. Exhibitors in the 1959 Annual Exposition of Science and Industry and descriptions of their exhibits.

The Directory

1. AAAS officers, staff, committees, for 1959.
2. Section committees and other AAAS Council members.
3. The 285 affiliated organizations.
4. Historical sketch and organization of the Association.
5. Complete roll of AAAS presidents and their fields.
6. Publications of the Association, including all symposium volumes.
7. AAAS Awards—including all past winners.
8. Future Meetings of the AAAS through 1963.
9. New and current activities of the AAAS.
10. Constitution and Bylaws.

Advance Registration

Advance registration has these decided advantages: 1) You avoid delay at the Registration Center upon arrival; 2) You receive the General Program in ample time to decide, unhurriedly, which events and sessions you particularly wish to attend; 3) Your name is posted in the Visible Directory as the Meeting opens.

The following coupon may be used both by advance registrants and by those who wish only the advance copy of the General Program; the Directory may also be ordered.

—THIS IS YOUR COUPON FOR AN ADVANCE COPY OF THE GENERAL PROGRAM— (AND FOR THE DIRECTORY, IF ALSO DESIRED)

- 1a. ☐ Enclosed is \$3.00 for my advance Registration Fee which brings me the General Program, Convention Badge, and all privileges of the Meeting.
- 1b. ☐ Enclosed is \$2.00 for only the General Program. (It is understood that, if I should attend the Meeting later, the Badge—necessary for the privileges of the Meeting—will be secured for \$1.00 more.)
- (Check 1a or 1b)
- 1c. ☐ Enclosed is \$1.00 for the Directory of AAAS Officers and Activities to be sent now.

Total amount enclosed \$

2. FULL NAME (Dr., Miss, etc.)
(Please print or typewrite) (Last) (First) (Initial)
ACADEMIC, PROFESSIONAL, OR
BUSINESS CONNECTION
4. OFFICE OR HOME ADDRESS
(For receipt of General Program and/or Directory)
5. FIELD OF INTEREST
6. CONVENTION ADDRESS
(May be added later, after arrival)

Please mail this Coupon and your check or money order for the total amount to the
AMERICAN ASSOCIATION FOR THE ADVANCEMENT OF SCIENCE
1515 Massachusetts Avenue, NW, Washington 5, D.C.

17-19. Northeast Electronics Research and Engineering Meeting, Boston, Mass. (Miss S. Whiteker, Inst. of Radio Engineers, 73 Tremont St., Boston, Mass.)

17-20. National Assoc. for Mental Health, annual, Philadelphia, Pa. (American Psychiatric Assoc., 1700 18 St., NW, Washington 9.)

18. Association for Psychiatric Treatment of Offenders, New York, N.Y. (M. Schmideberg, New York Acad. of Sciences, 2 E. 63 St., New York 21.)

19-21. Inter-Society Cytology Council, annual, Detroit, Mich. (P. A. Younge, ISCC, 1101 Beacon St., Brookline 46, Mass.)

20-21. American Mathematical Soc., Winston-Salem, N.C. (J. W. Green, Univ. of California, Los Angeles 24.)

20-21. Nuclear Fusion, symp., Austin, Tex. (Texas Symp. on Nuclear Fusion, P.O. Box 8005, University Station, Austin, Tex.)

22-24. American Soc. of Hematology, 2nd annual, St. Louis, Mo. (J. W. Rebuck, Henry Ford Hospital, Detroit, Mich.)

22-29. Pan American Child Cong., 11th, Bogotá, Colombia. (Office of Intern. Conferences, Department of State, Washington 25.)

23-24. Solid-State Techniques in Modern Instrumentation, symp., Philadelphia, Pa. (G. L. Eberly, 12 S. 12 St., Philadelphia 7.)

23-25. Fluid Dynamics (APS), Ann Arbor, Mich. (R. J. Emrich, Dept. of Physics, Lehigh Univ., Bethlehem, Pa.)

23-26. Technical European Conf. on Standards Applicable to Water (by invitation only), Copenhagen, Denmark. (World Health Organization, Regional Committee for Europe, 8 Scherfigsvej, Copenhagen.)

23-3. Inter-African Soils Conf., 3rd, Dalaba, Guinea. (Committee for Technical Cooperation in Africa South of the Sahara, Abbey House, 2-8 Victoria St., London, S.W.1, England.)

25. Association for the Advancement of Psychoanalysis, New York, N.Y. (New York Acad. of Medicine, 2 E. 103 St., New York, N.Y.)

26-27. Association for the Utilization of Atomic Energy in Ship-Building and Navigation, Hamburg, Germany. Gesellschaft für Kernenergieverwertung in Schiffbau m.b.H., Hamburg.)

26-28. Central Assoc. of Science and Mathematics Teachers, Chicago, Ill. (G. G. Mallinson, Western Michigan Univ., Kalamazoo.)

26-28. Ceylon Assoc. for the Advancement of Science, Colombo. (K. Arumugam and S. Wijesundera, General Secretaries, Univ. of Ceylon, Colombo 3.)

26-29. Legal and Administrative Problems of Peaceful Use of Nuclear Energy, intern. conf., Rio Piedras, Puerto Rico. (J. Mayda, Faculty of Law, Univ. of Puerto Rico, Rio Piedras.)

27-28. American Mathematical Soc., Detroit, Mich. (J. W. Green, Univ. of California, Los Angeles 24.)

27-28. American Physical Soc., Cleveland, Ohio. (K. K. Darrow, Columbia Univ., New York 27.)

27-28. American Soc. of Animal Production, Chicago, Ill. (H. H. Stonaker, Colorado State University, Fort Collins.)

(See issue of 18 September for comprehensive list)

New Products

The information reported here is obtained from manufacturers and from other sources considered to be reliable. Neither Science nor the writer assumes responsibility for the accuracy of the information. All inquiries concerning items listed should be addressed to Science, Room 740, 11 West 42 St., New York 36, N.Y. Include the name(s) of the manufacturer(s) and the department number(s).

■ **NEUTRON DETECTION FOILS** permit measurement of neutron-energy spectra as follows: gold and cobalt for thermal flux; Pu^{239} , energies greater than 1 kev; U^{235} , energies greater than 1.5 Mev; S^{32} , energies greater than 2.9 Mev; Al^{27} , using products of (n, p) reaction, energies greater than 5.3 Mev; Al^{27} and products of (n, α) reaction, energies greater than 8.6 Mev. (Aerojet-General Nucleonics, Dept. 99)

■ **SCRATCH INVESTIGATION MICROSCOPE** can be used to measure the depths of scratches on flat or curved surfaces. Range of measurement is 0.5 to 12×10^{-3} in. Oblique illumination of the object produces a magnification in depth of 4 times the lateral magnification. The instrument is portable and has battery-powered illumination. A camera attachment is available. (Hilger and Watts, Dept. 100)

■ **WATER DEMINERALIZER** of mixed-bed type connects directly to the tap to deliver water at full line pressure. Exchanger container may be removed and discarded or regenerated for reuse. Capacity is approximately 1500 grains of total solids as CaCO_3 . A conductivity meter is included. (Pfaudler Permutit, Dept. 101)

■ **NUMERAL INDICATOR TUBE** provides 1.5-in. high numerals said to be readable at distances up to 50 ft at a 160-deg angle with ambient illumination brighter than normal room illumination. The numeral to be illuminated is selected by grounding the appropriate connection. (Baird Atomic, Dept. 104)

■ **CHROMATOGRAM SCANNER** includes a synchronous stepping mechanism that advances the strip to be scanned in finite steps from 0 to $\frac{1}{4}$ in., as selected. At each step the strip is counted to preset time or preset count. A record of step number, elapsed time, count, and count rate is printed. (Tracerlab, Dept. 105)

■ **ANALOG COMPUTER** for gas chromatography is said to make possible the calculation of a complete analysis in less than 3 min with accuracy within ± 0.1 percent. The computer corrects peak areas or peak heights for differences in detector response and normalizes corrected areas to 100, thereby giving mole percentage or weight per-

Do you know that ...

SOLID MATTER is more than 99.99% empty space?
ELECTRONIC BRAINS solve all problems with only two figures: 0 and 1?
MAN has not five but at least ten physical senses?
THE MOON is better known to us than the interior of the earth?
COLOR exists only in the eye of the observer?
A STRAIGHT LINE is not always the shortest distance between two points?

You can find the answers to thousands of questions on science quickly and easily with

The Concise Dictionary of Science

by Frank Gaynor

An up-to-date dictionary providing concise definitions of terms and concepts pertaining to all fields of science. Full coverage is given to the newer sciences.

\$10.00

Philosophical Library, Publishers
15 East 40 Street, New York 16

STEREOSCOPIC MICROSCOPE for the budget-minded!

Designed and sturdily built for educational or industrial use, UNITRON Model MSL has the same optical and mechanical features found in higher priced instruments.

- Erect, three-dimensional image
- Coated optics and prisms
- Achromatic 2X paired objectives, paired wide field eyepieces
- Working distance 75mm.
- Vertical binocular body with interpupillary and diopter adjustments
- Total focusing excursion 70mm.
- Heavy base, removable glass stage plate, stage clips
- Choice of one magnification 10X, 20X or 30X, extra eyepieces available

Complete in fitted hardwood cabinet ...

FREE 10 DAY TRIAL

ALSO AVAILABLE

Model MSL-A with magnifications —
5X or 10X or 15X \$125.00
Extra eyepieces Pair \$19.50
Model LS Stereoscopic Microscope Lamp \$14.75



UNITRON

INSTRUMENT DIVISION of UNITED SCIENTIFIC CO.
204-206 MILK STREET • BOSTON 9, MASS.

Please rush to me, UNITRON's Microscope Catalog, 4-VI

Name _____
Company _____
Street _____
City _____ State _____

centage of each component. The three models available can handle 4-, 8-, or 12-component mixtures. (Podbielniak, Dept. 106)

■ **RESIDUAL CHLORINE ANALYZER** generates a current, by means of two electrodes made of dissimilar metals, proportional to the amount of free or total residual chlorine in a line. Electrodes are kept clean by small glass spheres that circulate across their surfaces. Residual chlorine is recorded in parts per million on standard recorder charts. (Fischer and Porter, Dept. 111)

■ **FILM CARD READER** for large-size documents copied on 35 mm microfilm features an 18-by-24-in. screen. A 500-watt lamp provides illumination. Adapters are available for roll film. (Eugene Dietzgen, Dept. 103)

■ **POWER SUPPLY** for special tubes furnishes 500 to 5000 volts at 0 to 10 ma. Output is positive, negative, or floating. Maximum ripple is 10 mv peak-to-peak. Voltage may be reset within ± 0.05 percent. (Alfred Electronics, Dept. 107)

■ **AIRBORNE AMPLIFIER** for thermocouple and strain-gage amplification and telemetering operations is designed to operate from 28-volt d-c or 6.3-volt,

400-cy/sec a-c. Input is ± 5 ma at over 20 kohm impedance. Gain is 400 to 1000, and bandwidth is 0 to 1000 cy/sec. Output impedance is less than 10 kohm. Operating temperature range is -55° to $+125^\circ\text{C}$; altitude range, up to 100,000 ft; vibration up to 10 grav at 2 kcy/sec. (Southwestern Industrial Electronics Co., Dept. 109)

■ **ACCESSORIES FOR TISSUE GRINDERS** include a motor, support, and cooling-chamber assembly. An adjustable-stroke mechanism permits up and down movement of the pestle within the homogenizer tube. The cooling chamber holds the tube firmly in an environment of crushed ice or other coolant. (Kontes Glass Co., Dept. 112)

■ **VAPOR-PROOF EMERGENCY SUIT** is opened and closed by a self-sealing slide fastener extending from the left knee, over the left shoulder, and around the head. A ratchet adjustment is provided for head-band size. All seams are cemented; no needle holes are present. Several types of respiratory apparatus are accommodated, and the suit may be furnished with a sound-powered communications system. The suit is designed to accommodate a man up to 6 ft 4 in. tall and weighing 250 lb. (Union Industrial Equipment Corp., Dept. 113)

■ **COMMUNICATION MONITOR** guards against failure of telephone communication lines. The instrument operates by sending an analog voice sample with the voice signal along a telephone line and comparing the two signals at the other end of the line being monitored. Differences in voltage between the compared signals indicate the degree of trouble. If service is unsatisfactory, the call is automatically switched to an alternate line. (International Telephone and Telegraph Corp., Dept. 110)

■ **RECTIFIER TEST SET** has forward-current ranges 0 to 1 and 0 to 5 amp. Reverse-voltage range is 0 to 1000 volts. Average forward voltage drop is measured from 0 to 1 volt, and average reverse current in four ranges from 0.1 to 100 ma. Four 1-percent meters are used for indication. (Wallson Associates, Dept. 118)

■ **DEGAUSSER** is designed to accommodate all sizes of instrumentation tapes and magnetic films. Tape is carried on a pushbutton-controlled, motor-driven turntable. Degaussing is accomplished in a predetermined 20-sec cycle. (Magnasync Manufacturing Co., Dept. 117)

JOSHUA STERN

National Bureau of Standards,
Washington, D.C.

Annual Review of MICROBIOLOGY

Volume 13

553 Pages

Editorial Committee: S. P. Chilton, C. E. Clifton,
C. A. Evans, R. E. Hungate, T. L. Jahn, W. W. Umbreit

Biosynthetic Aspects of
Metabolism

Aspects of Terminal
Respiration in
Microorganisms

Biology of Tetrahymena
Physiology and Develop-
ment of Lower Fungi
(Phycomycetes)

Antibiotics in Food
Preservation

Microbiological Aspects
of Tissue Culture

Newer Antibiotics

Plant Disease Resistance
Bacteria as Plant
Pathogens

Nematodes in Plant
Disease

Nonspecific Resistance
to Infections

Biosynthesis of Anti-
bodies, Facts and
Theories

Unicellular Clocks
Recombination in

Bacteriophage

New Viruses and Virus
Diseases of Man

Immunologic Tolerance
Complex Loci in

Microorganisms

Biochemical Cytology of
Microorganisms

Fermentative Metabolism
Bacterial Genetics (with

Particular Reference
to Genetic Transfer)

Sulphate Reduction by
Bacteria

\$7.00 postpaid (U.S.A.); \$7.50 postpaid (elsewhere)

ANNUAL REVIEWS, INC.

Grant Avenue, Palo Alto, California




Scientists the world over use
RODDER MODEL E because they
require a rugged **ULTRAMICROBALANCE**
with an **0.1 MICROGRAM SENSITIVITY**
and a 200 mg. load capacity.

For literature / testimonials write



**MICROTECH
SERVICES COMPANY**
775 Sunshine Drive
Los Altos, Calif., U.S.A.

HIGH SPEED DIRECT WRITING GALVANOMETER RECORDER



IMMEDIATELY VISIBLE TRACES WITHOUT PROCESSING

Available with 6 or 12 galvanometers and a variety of paper speeds. Trace velocities to 10,000 in./sec., frequency ranges to 3,000 c.p.s. Indispensable for static and dynamic analysis of strains, vibrations, pressures, etc. Daylight loading, records can be stored permanently. Superb quality, unmatched performance, moderate cost.

WRITE FOR FREE INFORMATION


C. H. STOELTING CO.

MULTI-CHANNEL OSCILLOGRAPHS • POLYGRAPHS, ALL KINDS
KYMOPHAPHS AND ACCESSORIES • DC SERVO-GRAPHIC RECORDER
CHOPPER AMPLIFIERS • STIMULATORS • STEREOTAXIC UNITS
MICRO-MANIPULATORS

424 NORTH HOMAN AVENUE, CHICAGO 24, ILLINOIS

FERMENTOR DRIVE ASSEMBLY FOR LABORATORY AND PILOT PLANT APPLICATION

- Aerobic or Anaerobic Fermentations • Tissue Cultures
- Other Metabolic Studies in Submerged Cultures



Holds 6 stainless steel FERMENTORS with pyrex jars of capacities 5, 7.5 or 14 liters each; removable for autoclaving. Leak-proof, non-freezing, ball-bearing agitator housings. Cool, quiet, continuous performance.

Stainless steel water baths, thermostatically controlled within $\pm 1/2^\circ\text{C}$. for temperatures up to 60°C . Twin, antifriction drives for wide range of agitation rates. Available with Automatic Anti-foam System. Single fermentor and 3-fermentor assemblies also available.

Write for Catalog F-S029



NEW BRUNSWICK SCIENTIFIC CO., INC.

PRECISION LABORATORY APPARATUS

P.O. BOX 606, NEW BRUNSWICK, NEW JERSEY

PERSONNEL PLACEMENT

CLASSIFIED: Positions Wanted, 25¢ per word, minimum charge \$4. Use of Box Number counts as 10 additional words. Payment in advance is required. Positions Open, \$33 per inch or fraction thereof. No charge for box number.

COPY for classified ads must reach SCIENCE 2 weeks before date of issue (Friday of every week).

DISPLAY: Rates listed below—no charge for Box Number. Monthly invoices will be sent on a charge account basis—provided that satisfactory credit is established.

Single insertion	\$33.00 per inch
4 times in 1 year	30.00 per inch
7 times in 1 year	28.00 per inch
13 times in 1 year	27.00 per inch
26 times in 1 year	26.00 per inch
52 times in 1 year	25.00 per inch

For **PROOFS** on display ads, copy must reach SCIENCE 4 weeks before date of issue (Friday of every week).

Replies to blind ads should be addressed as follows:

Box (give number)
Science
1515 Massachusetts Ave., N.W.
Washington 5, D.C.

POSITIONS WANTED

Enzymologist, young; publications in neurochemistry, marine toxins. Varied research experience, some teaching, seeks challenging research position. Box 224, SCIENCE. X

Ph.D., parasitology major, biochemistry and nutrition minors. Experience in zoology and public health teaching; research parasitologist of feed additive products. Desires research and development appointment. Medical Bureau, Burnice Larson, Director, 900 North Michigan, Chicago. X

POSITIONS OPEN

Biochemist, Ph.D., to actively integrate and expand clinical biochemistry section of outstanding specialized institution in southern California. Research potential. Salary \$12,000-\$14,000. Write to Box 219, SCIENCE. 10/2, 9

BIOCHEMIST GRADUATE (B.S. OR M.S.)

With some training in bacteriology to perform research in the isolation, identification and purification of products of microbial origin.

ATTRACTIVE SALARY

Employee benefits, 37 1/2 hours.

Send resume to

Ortho Pharmaceutical Corporation
Route 202
Raritan, New Jersey

Biochemist, M.D. or Ph.D., principal scientist for radioisotope research laboratory, 377 beds GMAS VA Hospital, Fort Howard, Md. Affiliated with Johns Hopkins and University of Maryland Medical Schools. Opportunity for basic as well as clinical investigation and medical application of isotopes. Experience required, knowledge of patient metabolism preferred. Starting salary \$8810 or \$10,130 per annum, depending on qualifications; liberal vacation and sick leave plan, career Civil Service. For further information, write or contact the Chief, Personnel Division, VA Hospital, 3900 Lock Raven Boulevard, Baltimore 18, Md. X

POSITIONS OPEN

Electron Microscopist, Ph.D. or equivalent. Research institute, New York. Modern well-equipped and well-staffed air-conditioned laboratory. Must have adequate training and experience in all phases of preparation of tissues and in handling machine. Apply Box 214, SCIENCE. 9/25; 10/2

Geneticist with strong biochemical background. Teaching with strong emphasis on research. Rank and salary dependent on training and experience although preference will be given to young Ph.D. with 2 to 3 years of postdoctorate experience. Appointment 1 July 1960. Good research facilities provided including nuclear reactor on campus. **Physiologist** with strong genetical interests may be considered. Applications with full particulars and references to Dr. J. A. Christensen, Saint Barnabas Medical Center, High St., Newark, N.J. 10/9, 16

Male or female, M.S. or Ph.D. Research laboratory electron microscopy of tissues and animal tissue cultures. Background in cytology or related field and experienced in operation of electron microscope. Send detailed résumé to Dr. J. A. Christensen, Saint Barnabas Medical Center, High St., Newark, N.J. 10/2

New World-Wide Graduate Award Directories for American scientists, teachers, librarians to subsidize their education and research. Stipends \$200—\$10,000. Volume I (1957), \$3; volume II (just published, no duplication), \$3; both volumes, \$5. Limited editions. Complete, specific information on 400 awards in United States and overseas in each volume. CRUSADE, Sci., Box 99, Station G, Brooklyn 22, N.Y. eow

Pharmacologist, Ph.D. with experience or training in drug evaluation and pharmaceutical research and development. Send résumé to Pharmaceutical Research Division, Norden Laboratories, Lincoln, Nebraska. X

POSITIONS OPEN

(a) **Organic Chemist**, Ph.D., 5 to 10 years' laboratory experience and **Bacteriologist**, master's or Ph.D. to head applications laboratory; opportunity, also for **Organic Chemist** to work on synthesizing of new organic compounds; research department; important company; \$12,000-\$15,000; East. (b) **Clinical Chemist**, Ph.D.; preferably with minimum of 5 years' recent experience as chemistry supervisor in hospital; 12-man group expanding facilities; Pacific Coast: \$8400-\$12,000. (c) **Organic Chemist** to become associated with two organic neurologists; Pacific Coast. (d) **Pharmaceutical Chemist** with product development and research experience; large city, Midwest; \$8000-\$10,000. (e) **Supervisor, Fermentation Process Development**; preferably biochemist or chemical engineer with administrative experience; important pharmaceutical company; college town near medical center; East. (f) **Biochemist** to join staff of biochemistry department of large pharmaceutical company; should have background in medical science and experience in clinical research; \$14,000; East. \$10-1 Medical Bureau, Burnside Larson, Director, 900 North Michigan Avenue, Chicago. X

(a) **Pharmacologist To Direct Clinical Research**; M.D. with previous pharmaceutical experience; able to organize and administer clinical research program; some travel; liaison with medical, related professions; young, rapidly growing mid-western concern; to \$18,000 or more. (b) **Bacteriologist**; M.S., Ph.D., to head section, increase work in serology, virology; 800-bed fully approved general hospital; prominent southern university city. (c) **Chemist**; B.S., M.S. to supervise busy routine laboratory, 350-bed university hospital; to \$7500; historic eastern city. (d) **Biochemist**; M.S., Ph.D. to do research in investigation allergenic extracts, fractionation, purification, characterization; to \$9000; northwestern branch allergenic research laboratories. Woodward Medical Bureau, Ann Woodward, Director, 185 North Wabash, Chicago. X

Research Director, Ph.D. Unusual opportunity for capable, energetic, and ambitious person to have full responsibility for laboratory in growing medium-size Chicago area pioneer manufacturer of ingredients used in pharmaceuticals, foods, and feeds. Laboratory functions include research, development, "trouble shooting," quality control. Industrial experience required in allied field. Excellent growth benefits. Box 210, SCIENCE. X

SCIENCE TEACHERS, LIBRARIANS, ADMINISTRATORS urgently needed for positions in many states and foreign lands. Monthly non-fee placement journal since 1952 gives complete job data, salaries. Members' qualifications and vacancies listed free. 1 issue, \$1. Yearly (12 issues) membership, \$5. CRUSADE, SCL, Box 99, Station G, Brooklyn 22, N.Y. ew

Translators (Russian and/or Chinese to English). Specialists in all branches of the physical sciences who can translate technical material on full- or part-time basis at home. Applicant must have degrees in their specific sciences, thorough familiarity with technical terminology involved, and native command of English. Write to Translation Editor, Consultants Bureau, Inc., 227 West 17 St., New York 11, N.Y. X

VIROLOGIST

IF you would like to work with an unusual group of biological scientists with unique responsibilities in data evaluation and experimental design. IF you like the academic atmosphere. IF you would enjoy working with both laboratory scientists and clinicians.

IF you are thoroughly trained in virology-immunology
YOU COULD BE THE SCIENTIST WE NEED.

To apply your talents to the many facets of an expanding program in vaccines and chemotherapy.

Send complete résumé to
L. H. WATSON
Employment Manager
SMITH KLINE
& FRENCH
LABORATORIES

1512 Spring Garden St.
Philadelphia 1, Pa.

The Market Place

BOOKS • SERVICES • SUPPLIES • EQUIPMENT

DISPLAY: Monthly invoices will be sent on a charge account basis—provided that satisfactory credit is established.

Single insertion	\$33.00 per inch
4 times in 1 year	30.00 per inch
7 times in 1 year	28.00 per inch
13 times in 1 year	27.00 per inch
26 times in 1 year	26.00 per inch
52 times in 1 year	25.00 per inch

For PROOFS on display ads, copy must reach SCIENCE 4 weeks before date of issue (Friday of every week).

BOOKS AND MAGAZINES

Your sets and files of scientific journals

are needed by our library and institutional customers. Please send us lists and description of periodical files you are willing to sell at high market prices. Write Dept. A35, CANNER'S, Inc., Boston 20, Massachusetts

WANTED TO PURCHASE against cash or in exchange:

- Annals of Missouri Botanical Garden, Vols. 1-39.
 - Botanical Gazette, Vols. 38-107.
 - Bull. of the Torrey Botanical Club, Vols. 1-54, Vols. 69-84.
 - Ecological monographs, Vol. 17 (1947): 2, 3 Vol. 18 (1948): 3.
 - Ecology, Vol. 3 (1922): 1.
- Proceedings of the National Academy of Sciences, Vol. 18, Vol. 20, Vols. 24-36. We offer:
- Annals of botany, N.S.: Vols. 2, 5, 11:41, 42, 12-15.
 - Bull. de la Société Botanique de France: Vol. 35 (1888), Vols. 97-101, Vols. 103-105 and a number of single issues.
 - Ecology: Vols. 18-23 and Vol. 28.
- Kindly address offers to: American Friends of the Hebrew University, 9 East 89th Street, New York 28, N.Y.

SUPPLIES AND EQUIPMENT

"From the hand of the veterinarian to research."



albino rats

CHARLES RIVER CD
(Caesarean derived)
CHARLES RIVER SD
(Sprague-Dawley descendants)
CHARLES RIVER W
(Wistar descendants)

HYPOPHYSECTOMIZED RATS

- Only Charles River CD animals used
- Rigidly controlled environment (same bldg.) birth to surgery.
- High speed surgery by graduate biologists.
- 10 years experience animal surgery.
- Overnight air service from Boston

THE CHARLES RIVER BREEDING LABS

Dept. B, 1093 Beacon Street, Brookline 46, Mass.
Henry L. Foster, D.V.M., President

FREE Subscription to AEROGRAF Research Notes, a technical publication on new developments in gas chromatography. We offer a complete selection of essential supplies and the Aerograph line of GLPC equipment.

Send for your free subscription today
WILKENS INSTRUMENT & RESEARCH INC.
Box 313, Walnut Creek, California
Yellowstone 5-1469

HYPOPHYSECTOMIZED RATS

Shipped to all points via Air Express
For further information write
HORMONE ASSAY LABORATORIES, Inc.
8159 South Spaulding Ave., Chicago 29, Ill.

Gas Chromatography

SUPPLIES AND EQUIPMENT

SWISS MICE



Send for booklet

TACONIC FARMS

GERMANTOWN NEW YORK

AMERICAN-LINCOLN

ANIMAL CAGES
and INCUBATORS



- American-Lincoln animal cages of all types ranging from single units to large four compartment sizes. Special cages, spare pans, racks, etc., built to order. Incubators for egg embryo and container work used by America's leading scientific institutions.

Write for New Catalog
AMERICAN-LINCOLN INCUBATOR CO.
New Brunswick, N.J. Dept. A5

MICE

C.F.1 I.C.R. DESCENDANTS
GREEN HILLS FARM

Box 381, Jeffersonville, New York
Tel.: 363

SPRAGUE-DAWLEY STRAIN* ALBINO RATS

* Descendants
Price lists sent on request

S & K Animal Co.
Box 479A, Jamesburg, N.J.

PROFESSIONAL SERVICES

LABORATORY SERVICES

for the FOOD, FEED, DRUG and CHEMICAL INDUSTRIES
Analyses, Biological Evaluation, Toxicity Studies, Insecticide Testing and Screening, Flavor Evaluation.

Project Research and Consultation

Write for Price Schedule
P. O. Box 2217 • Madison 1, Wis.

AUTHORS WANTED BY N.Y. PUBLISHER

New York, N.Y.—One of the nation's largest book publishers is seeking book-length manuscripts of all types—fiction, non-fiction, poetry. Special attention to new writers. For more information, send for booklet 28A—*it's free*. Vantage Press, 120 W. 31 St., New York 1. (Branches in Washington, D.C., Chicago, Hollywood, Calif.)

FOUNDED 1922

Food and Drug
Research Laboratories
INCORPORATED

Maurice Avenue at 58th Street
Maspeth 78, New York City
CABLE: FOODLABS TEL: TWINING 4-0800

IEC



**INTERNATIONAL
MODEL PR-2
REFRIGERATED
CENTRIFUGE**

... first choice for
blood fractionations and
similar low temperature
separations.

**HERE'S WHY MOST LABORATORY
DIRECTORS SPECIFY THE INTERNATIONAL PR-2
FOR LOW-TEMPERATURE SEPARATION**

... It offers positive temperature control with wide-range versatility for separations between -20° and $+10^{\circ}\text{C}$.

The temperature you set on the control panel is automatically held within $\pm 1^{\circ}\text{C}$. In fact, material temperature can be maintained at 0°C indefinitely even at speeds up to 19,000 rpm ... without subcooling rotor or chamber!

28 INTERCHANGEABLE HEADS ... horizontal, angle, high-speed, high-capacity and shaker ... provide accessory combinations to meet virtually all laboratory needs.

WIDE CAPACITY RANGE ... from 4 liters down to 7 ml. ... is unmatched in the refrigerated centrifuge class.


Write for Bulletin T.

INTERNATIONAL  EQUIPMENT CO.

1219 SOLDIERS FIELD ROAD, BOSTON 35, MASSACHUSETTS

Thomas
GAS METER
Dry Test • Portable
Laboratory Model

- for basal metabolism tests
- for estimating pulmonary function
- for measurement of air flow
in urea determinations



6023-B.

Direct reading, low resistance type. Capacity 35 liters of air per minute, with dial graduated in 100 ml divisions. Convenient for measurement of large volumes in respiratory studies. This compact meter, weighing 11 $\frac{3}{4}$ lbs. and equipped with handle, can be carried during exercise tests. Manometer and thermometer, connecting to inlet chamber, are included to permit correction to standard conditions.

Applications. Usable for basal metabolism tests by the Douglas bag method, as originally suggested by H. S. Newcomer, *Journal of Biological Chemistry*, Vol. XLVII, No. 3 (August, 1921); for estimating pulmonary function as described by Frederick C. Waring, Jr., in *The American Review of Tuberculosis*, Vol. 60, No. 2 (August, 1949), pp. 149-167; for measurement of air flow in urea determinations by the aeration method, etc.

Accuracy. Travel of large dial pointer is not perfectly uniform, so that reading may be in error by as much as ± 150 ml at 1/3rd maximum rated capacity. If used for other than large volume tests in which such an error would be negligible, it is recommended that the error be determined by displacing known volume of air with water or by reference to measuring instrument of known accuracy. Maximum accuracy is obtained by use at steady flow rate approaching the maximum capacity.

Meter Mechanism. Two-diaphragm, slide-valve, dry gas type, with soft, plastic bellows, 6 inches diameter, which drive the dial pointers with very little pressure. Residual

air volume approximately 5.5 liters. At maximum flow rate, pressure difference between inlet and outlet is approximately 10 mm of water (0.7 mm of mercury).

Meter Dial. Horizontal, uncovered, recessed in protective well on top of case. Diameter 6 inches, graduated to 10 liters in 100 ml divisions, with four small inset dials totaling to 100,000 liters. Pointers are exposed and can be turned directly to zero between tests without affecting accuracy of meter. Pointers of totalizing dials can be lifted off to facilitate resetting.

Manometer. U-type, for water filling. Attached to meter by threaded, swivel coupling. Adjustable scale is graduated on both edges to 50 mm above and below a central zero. Replaceable U-tube, made of Pyrex brand glass, 8 mm o.d. \times 1 mm wall, is mounted in end brackets of phenolic plastic and is armored by lateral strips of chromium plated brass.

Thermometer. Dial type, 2 inches diameter, range 0 to 50°C in 1/2° divisions.

Case. Of tin coated sheet steel, sweated seam construction, finished in gray enamel, with rubber feet and carrying handle. Ser-rated inlet and outlet tubes are 2 inches long and fit rubber tubing 1 inch bore. Outside dimensions, excluding fixtures, 7 $\frac{1}{4}$ \times 7 $\frac{1}{4}$ \times 12 $\frac{1}{2}$ inches high.

6023-B. Gas Meter, Dry Test, Portable, Laboratory Model, Thomas, as above described, complete with manometer, dial type thermometer, threaded cap for manometer port, and directions for use.....195.00



ARTHUR H. THOMAS COMPANY

More and more laboratories rely on Thomas

Laboratory Apparatus and Reagents

VINE ST. AT 3RD • PHILADELPHIA 5, PA.

